

SUBSTITUTED ACID DERIVATIVES USEFUL AS ANTIDIABETIC AND
ANTIOBESITY AGENTS AND METHOD


This is a continuation-in-part of U.S. application
5 Serial No. 09/664,598, filed September 18, 2000 which
application takes priority from U.S. provisional
application No. 60/155,400 filed September 22, 1999.

Field of the Invention

10 The present invention relates to novel substituted
acid derivatives which modulate blood glucose levels,
triglyceride levels, insulin levels and non-esterified
fatty acid (NEFA) levels, and thus are particularly
useful in the treatment of diabetes and obesity, and to a
15 method for treating diabetes, especially Type 2 diabetes,
as well as hyperglycemia, hyperinsulinemia,
hyperlipidemia, obesity, atherosclerosis and related
diseases employing such substituted acid derivatives
alone or in combination with another antidiabetic agent
20 and/or a hypolipidemic agent.

Description of the Invention

In accordance with the present invention,
substituted acid derivatives are provided which have the
25 structure I
I



wherein x is 1, 2, 3 or 4; m is 1 or 2; n is 1 or 2;

30 Q is C or N;
A is O or S;
Z is O or a bond;
R¹ is H or alkyl;
X is CH or N;



R^{2a} , R^{2b} and R^{2c} may be the same or different and are selected from H, alkyl, alkoxy, halogen, amino or substituted amino;

R³ is H, alkyl, arylalkyl, aryloxy carbonyl, alkyloxy carbonyl, alkynyloxy carbonyl, alkenyloxy carbonyl, aryl carbonyl, alkyl carbonyl, aryl, heteroaryl, alkyl(halo)aryloxy carbonyl, alkyloxy(halo)aryloxy carbonyl, cycloalkylaryloxy carbonyl, cycloalkyloxyaryloxy carbonyl, cycloheteroalkyl, heteroaryl carbonyl, heteroaryl-heteroarylalkyl, alkyl carbonylamino, aryl carbonylamino, heteroaryl carbonylamino, alkoxycarbonylamino, aryloxy carbonylamino, heteroaryloxy carbonylamino, heteroaryl-heteroaryl carbonyl, alkylsulfonyl, alkenylsulfonyl, heteroaryloxy carbonyl, cycloheteroalkyloxy carbonyl, heteroarylalkyl, aminocarbonyl, substituted aminocarbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylalkenyl, cycloheteroalkyl-heteroarylalkyl; hydroxyalkyl, alkoxy, alkoxyaryloxy carbonyl, arylalkyloxy carbonyl, alkylaryloxy carbonyl, arylheteroarylalkyl, arylalkylarylalkyl, aryloxyarylalkyl, haloalkoxyaryloxy carbonyl, alkoxycarbonylaryloxy carbonyl, aryloxyaryloxy carbonyl, arylsulfinylaryl carbonyl, arylthioaryl carbonyl, alkoxycarbonylaryloxy carbonyl, arylalkenyloxy carbonyl, heteroaryloxyarylalkyl, aryloxyaryl carbonyl, aryloxyarylalkyloxy carbonyl, arylalkenyloxy carbonyl, arylalkyl carbonyl, aryloxyalkyloxy carbonyl, arylalkylsulfonyl, arylthiocarbonyl, arylalkenylsulfonyl, heteroarylsulfonyl, arylsulfonyl, alkoxyarylalkyl, heteroarylalkoxycarbonyl, arylheteroarylalkyl, alkoxyaryl carbonyl, aryloxyheteroarylalkyl, heteroarylalkyloxyarylalkyl, arylarylalkyl, arylalkenylarylalkyl, arylalkoxyarylalkyl,

5 aminocarbonylarylalkyl;

10 structure $P(O)(OR^{4a})_2$, (where R^{4a} is H or a prodrug ester);


substituted with 1, 2 or 3 substituents;

15 thereof, with the proviso that

carbons in the normal chain.


20 have the structure

Ia



or


Ib




25

invention having the structure

5 IB














IC












$\begin{array}{c} \text{CH}_3 \quad \text{CH}_3 \\ \diagdown \quad \diagup \\ -\text{C}- \end{array}$, $(\text{CH}_2)_m$ is CH_2 , or $\begin{array}{c} \text{R}_a \\ | \\ -\text{CH}- \end{array}$ (where R_a is alkyl such as methyl, or alkenyl such as $-\text{CH}_2-\text{CH}=\text{CH}_2$ or $-\text{CH}_2-\text{C}(\text{CH}_3)=\text{CH}_2$), $(\text{CH}_2)_n$ is CH_2 , R^1 is lower alkyl,











- 4 -
























5

In addition, in accordance with the present invention, a method is provided for treating diabetes, especially Type 2 diabetes, and related diseases such as insulin resistance, hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, hyperlipidemia, obesity, hypertriglyceridemia, inflammation, Syndrome X, diabetic complications,

10

dysmetabolic syndrome, atherosclerosis, and related diseases wherein a therapeutically effective amount of a compound of structure I is administered to a human patient in need of treatment.

5 In addition, in accordance with the present invention, a method is provided for treating early malignant lesions (such as ductal carcinoma in situ of the breast and lobular carcinoma in situ of the breast), premalignant lesions (such as fibroadenoma of the breast and prostatic intraepithelial neoplasia (PIN),
10 liposarcomas and various other epithelial tumors (including breast, prostate, colon, ovarian, gastric and lung), irritable bowel syndrome, Crohn's disease, gastric ulceritis, and osteoporosis and proliferative diseases
15 such as psoriasis, wherein a therapeutically effective amount of a compound of structure I is administered to a human patient in need of treatment.

In addition, in accordance with the present invention, a method is provided for treating diabetes and related diseases as defined above and hereinafter, wherein a therapeutically effective amount of a combination of a compound of structure I and another type antidiabetic agent and/or a hypolipidemic agent, and/or lipid modulating agent and/or other type of therapeutic agent, is administered to a human patient in need of treatment.

In the above methods of the invention, the compound of structure I will be employed in a weight ratio to the antidiabetic agent (depending upon its mode of operation) within the range from about 0.01:1 to about 100:1, preferably from about 0.5:1 to about 10:1.

The conditions, diseases, and maladies collectively referenced to as "Syndrome X" or Dysmetabolic Syndrome are detailed in Johannsson J. *Clin. Endocrinol. Metab.*, 35 82, 727-734 (1997) and other publications.

The term "diabetes and related diseases" refers to Type II diabetes, Type I diabetes, impaired glucose

tolerance, obesity, hyperglycemia, Syndrome X, dysmetabolic syndrome, diabetic complications and hyperinsulinemia.

5 The conditions, diseases and maladies collectively referred to as "diabetic complications" include retinopathy, neuropathy and nephropathy, and other known complications of diabetes.

10 The term "other type(s) of therapeutic agents" as employed herein refers to one or more antidiabetic agents (other than compounds of formula I), one or more anti-obesity agents, and/or one or more lipid-lowering agents, one or more lipid modulating agents (including anti-atherosclerosis agents), and/or one or more antiplatelet agents, one or more agents for treating hypertension, one
15 or more anti-cancer drugs, one or more agents for treating arthritis, one or more anti-osteoporosis agents, one or more anti-obesity agents, one or more agents for treating immunomodulatory diseases, and/or one or more agents for treating anorexia nervosa.

20 The term "lipid-modulating" agent as employed herein refers to agents which lower LDL and/or raise HDL and/or lower triglycerides and/or lower total cholesterol and/or other known mechanisms for therapeutically treating lipid disorders.

25

Detailed Description of the Invention

The compounds of the formula I of the present invention may be prepared according to the following
30 general synthetic schemes, as well as relevant published literature procedures that are used by one skilled in the art. Exemplary reagents and procedures for these

5 Protecting Groups in Organic Synthesis, 3rd Edition, 1999
[Wiley]).

10 methyl-oxazole-4-ethanol) is coupled with a hydroxy aryl-
or heteroaryl- aldehyde III (preferably 3- or 4-
hydroxybenzaldehyde) under standard Mitsunobu reaction
conditions (e.g. Mitsunobu, O., *Synthesis*, **1981**, 1).
The resulting aldehyde IV is then subjected to reductive
15 amination using procedures known in the literature (e.g.
Abdel-Magid et al, *J. Org. Chem.* **1996**, 61, 3849) with an
 α -amino ester hydrochloride V. PG in Scheme 1 denotes a
preferred carboxylic acid protecting group, such as a
methyl or tert-butyl ester. The resulting secondary
20 amino-ester VI is then subjected to a second reductive
amination using methods known in the literature (e.g.
Abdel-Magid et al, *J. Org. Chem.* **1996**, 61, 3849) with an
R^{3a} aldehyde VII. Final deprotection of the carboxylic
acid ester under standard conditions known in the
25 literature (Greene) utilizing basic conditions (for
methyl esters) or acidic conditions (for tert-butyl
esters) then furnishes the desired amino acid products
ID.

30 Scheme 1A. The alcohol II ($R^5(CH_2)_xOH$) (of which the most favored is 2-[2-phenyl-5-methyl-oxazole-4-yl]-ethanol) is treated with methanesulfonyl chloride to give the corresponding mesylate VIII. The mesylate is then alkylated under standard basic conditions with a
35 hydroxyaryl or hydroxyheteroaryl aldehyde III to furnish the aldehyde IV.

Alternatively, as shown in Scheme 3, the tertiary amino acids IF may also be obtained by alkylation of the secondary amino-ester VI with an alkylating agent IX (with an appropriate leaving group (LG) such as halide, mesylate, or tosylate) under standard conditions known in the art followed again by standard deprotection of the carboxylic acid ester X to provide the amino acids IF.

For further substituted amino acids, a general synthetic scheme is shown in Scheme 5. Reductive amination of an appropriate amine XIII with an aryl or heteroaryl aldehyde XIV under standard conditions furnishes the corresponding secondary amine XV, which is then reacted with a halide-ester XVI (e.g. tert-butyl bromoacetate) to furnish the corresponding α -amino ester XVII. This amine-ester XVII is then deprotected under standard conditions to provide the desired amino acid analogs IF.

- 19 -

protected halide-phosphonate XVIIA to provide the corresponding aminophosphonate ester XVIIIA, which is then deprotected under standard conditions (Greene & Wuts) to furnish the amino phosphonic acid IFA. Scheme 5b illustrates the synthesis of the aminophosphinic acids IFB, which again involves the reaction of an appropriately protected halide-phosphinate ester XVIB with the secondary amine XV. Deprotection of the resulting aminophosphinate ester then provides the phosphinic acid IFB.

An alternative to the sequence in Scheme 5 is shown in Scheme 6. A hydroxyaryl or heteroaryl amine XVIII is selectively protected on nitrogen to provide protected amine XIX. A preferred $R^5(CH_2)_nOH$ (II) is then reacted with XIX under Mitsunobu conditions to provide the corresponding ether, followed by deprotection of the amine, to form the free amine XX. The free amine XX is then activated with a standard activating group (2,4-dinitrobenzenesulfonamide; T. Fukuyama et al, *Tetrahedron Lett.* **1997**, 38, 5831) and is then treated with an α -halo ester XVI as in Scheme 5. The 2,4 dinitrobenzenesulfonamide XXI is deprotected under literature conditions (T. Fukuyama et al, *Tetrahedron Lett.*, **1997**, 38, 5831) to furnish a secondary α -amino-ester XXII which is then subjected to a reductive amination with an R^{3a} aldehyde XI followed by deprotection of the ester X to furnish the desired analogs IF.

Scheme 7 describes an alternative general route to the amino acid analogs IG. A hydroxyaryl or heteroaryl aldehyde III is subjected to the usual reductive amination conditions with an appropriate amine-ester hydrochloride V. The resulting secondary amine-ester XXIII is functionalized, in this case by a second reductive amination with an R^{3a} aldehyde VII to furnish the corresponding hydroxy tertiary amine-ester XXIV. This can now undergo a Mitsunobu reaction with a preferred alcohol II ($R^5(CH_2)_nOH$) which followed by



10

20

30

Scheme 11 illustrates the synthesis of the carbamate-acid analogs IK. The secondary amine-ester XXII can be reacted with appropriate chloroformates XXXV under standard literature conditions (optimally in CH_2Cl_2 or CHCl_3 in the presence of a base such as Et_3N) to furnish the corresponding carbamate-esters. The requisite analogs IK are then obtained after deprotection of the carbamate-ester. Alternatively, the secondary amine-ester XXII can be reacted with phosgene to generate the corresponding carbamyl chloride XXXVI. This carbamyl chloride intermediate XXXVI can be reacted with $\text{R}^{3a}\text{-OH}$ (XXXVII) (optimally substituted phenols) to afford the corresponding carbamate-acids IK after deprotection.

Scheme 12 illustrates the further functionalization of aryl carbamate-acid analogs IK. The secondary amine-ester XXII is reacted with an aryl chloroformate XXXVIII (containing a protected hydroxyl group) to form XXXIX. The hydroxyl group is then selectively deprotected in the presence of the ester functionality to provide XL, then alkylated with an appropriate $\text{R}^6\text{-LG}$ (XLI) (where LG is halide, mesylate or tosylate, and R^6 is most preferably $\text{CHF}_2\text{-}$, or $\text{CH}_3\text{CH}_2\text{-}$) in the presence of base. Deprotection of the ester then furnishes the desired carbamate-acid analogs IL.

The secondary amine-ester XXIIA can be functionalized with substituted aryl or aliphatic carboxylic acids XLII, under standard peptide coupling conditions, as illustrated in Scheme 13. The amide bond-forming reactions are conducted under standard peptide coupling procedures known in the art. Optimally, the reaction is conducted in a solvent such as DMF at 0°C to RT using 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDAC or EDCI or WSC), 1-hydroxybenzotriazole (HOBT) or 1-hydroxy-7-azabenzotriazole (HOAT) and a base, for example Hunig's base (diisopropylethylamine), N-methylmorpholine or triethylamine. Deprotection of the amide-ester then furnishes the desired amide-acid analogs IM.

The secondary amine-ester XXIIA can also be reacted with aliphatic or aryl isocyanates XLIII to provide the corresponding urea-esters. Deprotection of the urea-ester provides the desired urea-acid analogs IN, as shown in Scheme 14. Alternatively, as shown in Scheme 15, the carbamyl chloride intermediate XXXVI described in Scheme 11 can be reacted with appropriate aliphatic or aryl amines XLIV in the presence of a tertiary amine (e.g. Et₃N) to furnish tri- or tetrasubstituted urea-acid analogs IO or IP after deprotection of the ester.


The secondary amine-ester XXIIA can also be reacted with appropriate sulfonyl chlorides XLVI under standard literature conditions (optimally in the presence of a base such as pyridine, either neat or using chloroform as a co-solvent), followed by deprotection, to provide the corresponding sulfonamide-acids IQ, as shown in Scheme 16.

Replacement of the carboxylic acid functionality in these analogs with tetrazole can be achieved as shown in Scheme 17. An acid analog IK is coupled with an amine (containing an appropriate tetrazole protecting group) XLVII (preferably 3-amino propionitrile) under standard peptide coupling conditions. The resulting secondary amide XLVIII is then subjected to a Mitsunobu reaction under standard conditions with trimethylsilyl azide (TMSN₃) to form the protected tetrazole XLIX. Deprotection of the cyanoethyl group is achieved preferentially in the presence of base to generate the desired free tetrazole analog IR.

Scheme 18 describes a general synthesis of the hydrazide-acid analogs IS. A substituted aryl carboxylic acid 1 is treated with methanesulfonyl chloride in the presence of base; the intermediate is then reacted with a protected hydrazine-ester VA to give the corresponding acylated hydrazine 1a (ref: *Synthesis*, 1989, 745-747). The acylhydrazine 1a is coupled with an appropriately substituted aryl aldehyde IV under reductive amination

Deprotection of the ester 3 then furnishes the hydrazide-acid analogs 1S.

15 The different approaches to the preparation of the
α-alkylbenzyl amino acid and carbamate-acid analogs IT
and IU are exemplified in the following synthetic
schemes. In Scheme 20 an appropriately substituted aryl
aldehyde IV is treated with a suitable organometallic
20 reagent (e.g. a Grignard reagent $R^{10}MgBr$) under standard
conditions to give the corresponding secondary alcohol,
which is then oxidized under standard conditions (e.g.
Swern oxidation with $(COCl)_2/DMSO/Et_3N$ or using pyridinium
chlorochromate) to give the corresponding ketone 5.
25 Reductive amination of the ketone 5 with an appropriately
substituted amino-ester 6 provides the corresponding α-
alkylbenzyl amino-ester 7. It will be understood that in

the amino ester 6, the moiety  does not necessarily represent two repeating units.

30 Acylation of amino-ester 7 with an appropriately substituted aryl or heteroaryl chloroformate XXXV followed by deprotection provides the racemic carbamate-acid analogs IT. Reductive amination of alkylbenzyl amino-ester 7 with aryl aldehyde VII followed by
35 deprotection provides the racemic amino-acid analogs IU.

Alternatively, as shown in Scheme 21, asymmetric reduction (e.g. using the Corey oxazaborolidine reduction protocol; review: E. J. Corey & C. Helal, *Angew. Chem. Int. Ed. Engl.*, 1998, 37, 1986-2012.) of the aryl-ketone 5 provides each of the desired enantiomeric alcohols 8 (although only one enantiomer is represented in the scheme). Treatment of the chiral alcohol 8 with azide in a Mitsunobu-like reaction (ref: A. S. Thompson et. al., *J. Org. Chem.* 1993, 58, 5886-5888) gives the corresponding chiral azide (with inverted stereochemistry from the starting alcohol). The azide is then converted to the amine 9 by standard reduction methods (e.g. hydrogenation or $\text{Ph}_3\text{P}/\text{THF}/\text{H}_2\text{O}$). Treatment of the chiral amine 9 with an ester XVIA (containing an appropriate leaving group) provides the secondary amino-ester 10. Acylation of amino-ester 10 with an aryl or heteroaryl chloroformate XXXV followed by deprotection provides the chiral carbamate-acid analogs ITa (which may be either enantiomer depending upon the stereochemistry of 8). Reductive amination of alkyl amino-ester 10 with aryl aldehydes VII followed by deprotection provides the chiral amino-acid analogs IUa (which may be either enantiomer depending upon the stereochemistry of 8).

An alternative to Scheme 21 is shown in Scheme 22. An appropriately protected oxyaryl ketone 11 undergoes asymmetric reduction to give the chiral alcohol 12. This is converted to the chiral amine 13 via the identical sequence as in Scheme 21 (via the chiral azide). Treatment of the chiral amine 13 with an ester XVIA (LG = halogen or mesylate) gives the corresponding secondary amino-ester 14. Acylation of 14 with an aryl or heteroaryl chloroformate XXXV provides the corresponding carbamate-ester. Selective deprotection furnishes the free phenol carbamate-ester 15. Alkylation of the phenol 15 with a halide or mesylate VIII followed by deprotection provides the carbamate-acid analogs ITa. An analogous sequence (involving reductive amination of the

secondary amino-ester 14 with an aryl or heteroaryl aldehyde VII, then selective deprotection, alkylation with VIII and a final deprotection) provides the amino acid analogs IUa.

5 It will be appreciated that either the (R)- or (S)- enantiomer of ITa or IUa may be synthesized in Schemes 21 and 22, depending upon the chirality of the reducing agent employed.

 A fourth synthetic sequence is shown in Scheme 23.
10 The substituted aldehyde IV is condensed with an amino- ester hydrochloride 6 to give the corresponding imine 16, which is then treated in situ with an appropriately substituted allylic halide 17 in the presence of indium metal (reference: Loh, T.-P., et al., *Tetrahedron Lett.*,
15 1997, 38, 865-868) to give the α -allyl benzyl amino-ester 18. Acylation of amine 18 with an aryl or heteroaryl chloroformate XXXV followed by deprotection provides the carbamate-acid analogs Iv. Reductive amination of alkyl amino-ester 18 with an aryl or heteroaryl aldehyde VII
20 followed by deprotection provides the amino-acid analogs IW.

 Scheme 24 shows the preparation of the required intermediate 2-aryl-5-methyl-oxazol-4-yl methyl chloride 21 (following the general procedure described in Malamas,
25 M. S., et al, *J. Med. Chem.*, 1996, 39, 237-245). A substituted aryl aldehyde 19 is condensed with butane- 2,3-dione mono-oxime under acidic conditions to give the corresponding oxazole N-oxide 20. Deoxygenation of the oxazole N-oxide 20 with concomitant chlorination
30 furnishes the desired chloromethyl aryl-oxazoles 21. Hydrolysis of chloromethyl oxazole 21 under basic conditions furnishes the corresponding oxazole-methanol 22. Oxidation of alcohol 22 to the corresponding aldehyde is followed by conversion to the corresponding
35 dibromoalkene 23 (e.g. $\text{Ph}_3\text{P/CBr}_4$). The dibromide 23 is converted to the corresponding alkynyl-lithium species (using an organolithium reagent such as n-BuLi), which

can be reacted in situ with an appropriate electrophile such as formaldehyde to give the corresponding acetylenic alcohol (ref: Corey, E. J., et al., *Tetrahedron Lett.* 1972, 3769, or Gangakhedkar, K. K., *Synth. Commun.* 1996, 26, 1887-1896). This alcohol can then be converted to the corresponding mesylate 24 and alkylated with an appropriate phenol 25 to provide analogs IX. Further stereoselective reduction (e.g. H₂/Lindlar's catalyst) provides the E- or Z- alkenyl analogs IY.

10 Scheme 25 describes a general synthesis of the amino-benzoxazole analogs IZ (general ref: Sato, Y., et al, *J. Med. Chem.* 1998, 41, 3015-3021). An appropriately substituted ortho-aminophenol 26 is treated with CS₂ in the presence of base to furnish the corresponding
15 mercapto-benzoxazole 27. Treatment of this thiol 27 with an appropriate chlorinating agent (e.g. PCl₅) provides the key intermediate chlorobenzoxazole 28, which is reacted with the secondary amino-ester VI to furnish, after deprotection, the amino benzoxazole-acid analogs IZ.

20 The thiazole analogs IZa were synthesized according to the general synthetic route outlined in Scheme 26 (ref. Collins, J. L., et al., *J. Med. Chem.* 1998, 41, 5037). The secondary amino-ester XXIII is reacted with an aryl or heteroaryl chloroformate XXXV in the presence
25 of an appropriate base (e.g. pyridine or triethylamine) to furnish the corresponding hydroxyaryl carbamate-ester 29. The hydroxyaryl ester 29 is then reacted with an appropriately substituted α -bromo vinyl ketone 29a (for S₃ = CH₃, e.g. Weyerstahl, P., et. al., *Flavour Fragr. J.*,
30 1998, 13, 177 or Sokolov, N. A., et al., *Zh. Org. Khim.*, 1980, 16, 281-283) in the presence of an appropriate base (e.g. K₂CO₃) to give the corresponding Michael reaction adduct, the α -bromoketone carbamate-ester 30. The α -bromoketone 30 is then subjected to a condensation
35 reaction with an appropriately substituted aryl amide 31 (A = O) or aryl thioamide 31 (A = S) to furnish either the corresponding oxazole (from the amide) or the

thiazole (from the thioamide) (ref: Malamas, M. S., et al, *J. Med. Chem.*, 1996, 39, 237-245). Finally, deprotection of esters 31 then provides the substituted oxazole and thiazole carbamate acid analogs IZa.

5 It will be appreciated that in the following
schemes where the carbamate-acid analogs are prepared,
the corresponding amino acid analogs may also be prepared
by replacing the chloroformate reaction with an aldehyde
in a reductive amination reaction (as in Scheme 20 with
10 intermediate amine 7).

Scheme 27 describes a general synthesis of the acids IZb and IZc. A halo-substituted aryl aldehyde 32 (preferably iodide or bromide) is subjected to reductive amination using procedures known in the literature (e.g. Abdel-Magid et al, *J. Org. Chem.* 1996, 61, 3849) with an α -amino acid ester hydrochloride V. The resulting secondary amino-ester 33 is then reacted with an aryl or heteroaryl chloroformate XXXV in the presence of an appropriate base (e.g. pyridine or triethylamine) to furnish the corresponding halo-aryl carbamate-ester 34. Aryl halide 34 is then reacted with an appropriate aryl- or heteroaryl-substituted acetylene 35 (the preferred acetylene being 5-phenyl-2-methyl-oxazol-4-yl-methylacetylene) in the presence of an appropriate palladium catalyst (e.g. $(\text{Ph}_3\text{P})_2\text{PdCl}_2$) and a copper (I) salt (e.g. CuI) in a Sonogashira coupling reaction (ref: *Organocopper Reagents, a Practical Approach*, R. J. K. Taylor, Ed., Chapter 10, pp 217-236, Campbell, I. B., Oxford University Press, 1994) to furnish the key intermediate, arylacetylene carbamate ester 36.

The arylacetylene ester 36 is deprotected to provide the corresponding arylacetylene acid analogs 1Zb. The acetylene moiety of 36 can be reduced by standard methods (e.g. hydrogenation, ref: M. Hudlicky, 35 Reductions in Organic Chemistry, 2nd Edition, ACS, 1996, Chapter 1) to furnish the corresponding fully saturated

alkyl aryl carbamate ester, which is then deprotected to give the alkyl aryl carbamate acid analogs IZc.

Stereoselective reduction of the acetylene ester 36 by standard methods (e.g. Lindlar's catalyst; ref: 5 Preparation of Alkenes, A Practical Approach, J. J. Williams, Ed., Chapter 6, pp 117-136, Oxford University Press, 1996) can be achieved to provide the corresponding cis-alkenyl aryl carbamate-ester, which is then deprotected to furnish the Z-alkenyl aryl carbamate acid 10 analogs IZd (Scheme 28). Alternatively, this sequence can be reversed, i.e. the initial step being the deprotection of acetylenic ester 36 to the acetylenic acid, followed by stereoselective reduction of the acetylene moiety to provide the Z-alkene-acid analogs 15 IZd.

The corresponding trans-alkenyl aryl carbamate acids IZe can be synthesized according to the general route in Scheme 29. An aryl- or heteroaryl-acetylene 35 (the preferred moiety again being 5-phenyl-2-methyl-oxazol-4-yl-methylacetylene) is halogenated under 20 standard conditions (ref: Boden, C. D. J. et al., *J. Chem. Soc. Perkin Trans. I*, 1996, 2417; or Lu, W. et. al., *Tetrahedron Lett.* 1998, 39, 9521) to give the corresponding halo-acetylene, which is then converted to 25 the corresponding trans-alkenyl stannane 37 (ref: Boden, C. D. J., *J. Chem. Soc., Perkin Trans. I*, 1996, 2417). This aryl- or heteroaryl-substituted trans-alkenyl stannane 37 is then coupled with the halo-aryl carbamate ester 34 under standard Stille coupling conditions (ref: 30 Farina, V. et. al., "The Stille Reaction", *Organic Reactions*, 1997, 50, 1) to furnish the corresponding trans-alkenyl aryl carbamate ester 38. This carbamate-ester is then deprotected under standard conditions to give the desired trans-alkenyl aryl carbamate acid 35 analogs IZe.


The corresponding cyclopropyl analogs IZf and IZg are synthesized according to Scheme 30. For the cis- or

(Z-) cyclopropyl analogs, stereoselective reduction (H_2 /Lindlar's catalyst) of the alkynyl moiety of intermediate alkynyl ester 36 (as for analogs IZd), followed by cyclopropanation under standard conditions (Zhao, Y., et al, *J. Org. Chem.* 1995, 60, 5236-5242) and finally deprotection provides the cis-cyclopropyl carbamate-acid analogs IZf. For the trans-cyclopropyl analogs IF, analogous cyclopropanation of the E-alkene moiety of intermediate 38 followed by deprotection provides the trans-cyclopropyl carbamate-acid analogs IZg.


A preferred alternative asymmetric synthesis of ITa (Scheme 21) is shown in Scheme 31. Protection of a chiral amine 39 (with the phenol protected), preferably as a carbamate, provides intermediate 40. Removal of the phenolic protecting group of 40 provides the free phenol 41. Alkylation of phenol 41 with the mesylate VIII furnishes the protected amine 42. Deprotection of this amine then furnishes the key intermediate secondary amino-ester 9, which is then carried on to analogs ITa and IUa according to Scheme 21.

A preferred asymmetric synthesis of analogs IIA is shown in Scheme 32. The aldehyde IV is subjected to standard Wittig reaction conditions (ref: Preparation of Alkenes, a Practical Approach, J. J. Williams, Ed., Chapter 2, pp 19-58) to furnish the alkene 43. Asymmetric aminohydroxylation according to known literature procedures (ref: O'Brien, P., *Angew. Chem. Int. Ed.*, 1999, 38, 326 and Reddy, K. L., and Sharpless, K. B., *J. Am. Chem. Soc.*, 1998, 120, 1207) furnishes the desired amino-alcohol 44 as a single enantiomer. Selective protection of the amine provides the alcohol 45. Alcohol 45 is then converted to the intermediate 46, which contains a suitable leaving group (either a halide or a mesylate) for the subsequent cuprate reaction. Reaction of an appropriate higher-order cuprate (ref: L. A. Paquette, Ed., *Organic Reactions*, 1992, Vol. 41, J. Wiley & Sons) with the protected amine substrate 46


5



In this and the following Reaction Schemes:




Alternative Scheme 1A for Preparing Aldehyde IV




SCHEME 1A






Scheme 4



Scheme 5




- 35 -



Scheme 5b









S_1 = H, alkyl, halo, alkoxy, alkylthio, alkylamino, aryloxy, aryl, heteroaryl, alkoxy carbonyl, alkylaminocarbonyl
 S_2 = H, alkyl, halo, alkoxy, alkylthio, alkylamino, aryloxy, aryl, heteroaryl, alkoxy carbonyl, alkylaminocarbonyl

Scheme 8




Scheme 9





Scheme 11






- 43 -











Scheme 21




Scheme 22






Scheme 24




- 52 -






Scheme 27




Scheme 28



Scheme 29








Scheme 32

Unless otherwise indicated, the term "lower alkyl", "alkyl" or "alk" as employed herein alone or as part of another group includes both straight and branched chain hydrocarbons, containing 1 to 20 carbons, preferably 1 to 10 carbons, more preferably 1 to 8 carbons, in the normal chain, and may optionally include an oxygen or nitrogen in the normal chain, such as methyl, ethyl, propyl, isopropyl, butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl, dodecyl, the various branched chain isomers thereof, and the like as well as such groups including 1 to 4 substituents such as halo, for example F, Br, Cl or I or CF₃, alkoxy, aryl, aryloxy, aryl(aryl) or diaryl, arylalkyl, arylalkyloxy, alkenyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkyloxy, amino, hydroxy, hydroxyalkyl, acyl, heteroaryl, heteroaryloxy, cycloheteroalkyl, arylheteroaryl, arylalkoxycarbonyl, heteroarylalkyl, heteroarylalkoxy, aryloxyalkyl, aryloxyaryl, alkylamido, alkanoylamino, arylcarbonylamino, nitro, cyano, thiol, haloalkyl, trihaloalkyl and/or alkylthio and/or any of the R³ groups.


Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 or 2 double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 10 carbons, forming the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl,





any of which groups may be optionally substituted with 1 to 4 substituents such as halogen, alkyl, alkoxy, hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the substituents for alkyl.

The term "cycloalkenyl" as employed herein alone or as part of another group refers to cyclic hydrocarbons containing 3 to 12 carbons, preferably 5 to 10 carbons and 1 or 2 double bonds. Exemplary cycloalkenyl groups include cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, cyclohexadienyl, and cycloheptadienyl, which may be optionally substituted as defined for cycloalkyl.

The term "cycloalkylene" as employed herein refers to a "cycloalkyl" group which includes free bonds and thus is a linking group such as



,  and the like, and may optionally be substituted as defined above for "cycloalkyl".

The term "alkanoyl" as used herein alone or as part of another group refers to alkyl linked to a carbonyl group.

Unless otherwise indicated, the term "lower alkenyl" or "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 1 to 8 carbons in the normal chain, which include one to six double bonds in the normal chain, and may optionally include an oxygen or nitrogen in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-

5

10

30


35





Where alkenyl groups as defined above and alkynyl groups as defined above, respectively, have single bonds for attachment at two different carbon atoms, they are

termed "alkenylene groups" and "alkynylene groups", respectively, and may optionally be substituted as defined above for "alkenyl" and "alkynyl".

(CH₂)_x, (CH₂)_m, (CH₂)_n or (CH₂)_y includes alkylene, allenyl, alkenylene or alkynylene groups, as defined herein, each of which may optionally include an oxygen or nitrogen in the normal chain, which may optionally include 1, 2, or 3 substituents which include alkyl, alkenyl, halogen, cyano, hydroxy, alkoxy, amino, thioalkyl, keto, C₃-C₆ cycloalkyl, alkylcarbonylamino or alkylcarbonyloxy; the alkyl substituent may be an alkylene moiety of 1 to 4 carbons which may be attached to one or two carbons in the (CH₂)_x or (CH₂)_m or (CH₂)_n group to form a cycloalkyl group therewith.

15 Examples of $(\text{CH}_2)_x$, $(\text{CH}_2)_m$, $(\text{CH}_2)_n$, $(\text{CH}_2)_y$,
alkylene, alkenylene and alkynylene include







10 The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine as well as CF₃, with chlorine or fluorine being preferred.

15 The term "metal ion" refers to alkali metal ions such as sodium, potassium or lithium and alkaline earth metal ions such as magnesium and calcium, as well as zinc and aluminum.

Unless otherwise indicated, the term "aryl" or the group



20 where Q is C, as employed herein alone or as part of another group refers to monocyclic and bicyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including 1-naphthyl and 2-naphthyl) and may optionally include one to three
25 additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings for example



5

and may be optionally substituted through available carbon atoms with 1, 2, or 3 groups selected from hydrogen, halo, haloalkyl, alkyl, haloalkyl, alkoxy,
 10 haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkyl-alkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, alkoxycarbonyl, arylcarbonyl, arylalkenyl, aminocarbonylaryl, arylthio,
 15 arylsulfinyl, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio,
 20 arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino,
 25 arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfonaminocarbonyl and/or any of the substituents for alkyl set out herein.

Unless otherwise indicated, the term "lower alkoxy", "alkoxy", "aryloxy" or "aralkoxy" as employed
 30 herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to an oxygen atom.

Unless otherwise indicated, the term "substituted amino" as employed herein alone or as part of another group refers to amino substituted with one or two substituents, which may be the same or different, such as
5 alkyl, aryl, arylalkyl, heteroaryl, heteroarylalkyl, cycloheteroalkyl, cycloheteroalkylalkyl, cycloalkyl, cycloalkylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl or thioalkyl. These substituents may be further substituted with a carboxylic acid and/or any of the substituents for
10 alkyl as set out above. In addition, the amino substituents may be taken together with the nitrogen atom to which they are attached to form 1-pyrrolidinyl, 1-piperidinyl, 1-azepinyl, 4-morpholinyl, 4-thiamorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-
15 arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, 1-pyrrolidinyl, 1-piperidinyl, or 1-azepinyl, optionally substituted with alkyl, alkoxy, alkylthio, halo, trifluoromethyl or hydroxy.

Unless otherwise indicated, the term "lower alkylthio", "alkylthio", "arylthio" or "aralkylthio" as
20 alkylthio", "alkylthio", "arylthio" or "aralkylthio" as employed herein alone or as part of another group includes any of the above alkyl, aralkyl or aryl groups linked to a sulfur atom.

Unless otherwise indicated, the term "lower alkylamino", "alkylamino", "arylamino", or
25 "arylalkylamino" as employed herein alone or as part of another group includes any of the above alkyl, aryl or arylalkyl groups linked to a nitrogen atom.

Unless otherwise indicated, the term "acyl" as
30 employed herein by itself or part of another group, as defined herein, refers to an organic radical linked to a carbonyl $\left(\begin{smallmatrix} \text{O} \\ \parallel \\ \text{C} \end{smallmatrix} \right)$ group; examples of acyl groups include any of the R³ groups attached to a carbonyl, such as alkanoyl, alkenoyl, aroyl, aralkanoyl, heteroaroyl,
35 cycloalkanoyl, cycloheteroalkanoyl and the like.

Unless otherwise indicated, the term "cycloheteroalkyl" as used herein alone or as part of

5




20

25



25

possible N-oxides. The heteroaryl group may optionally include 1 to 4 substituents such as any of the the substituents for alkyl or aryl set out above. Examples of heteroaryl groups include the following:



and the like.

The term "cycloheteroalkylalkyl" as used herein
15 alone or as part of another group refers to
cycloheteroalkyl groups as defined above linked through a
C atom or heteroatom to a (CH₂)_p chain.

The term "heteroarylalkyl" or "heteroarylalkenyl" as used herein alone or as part of another group refers to a heteroaryl group as defined above linked through a C atom or heteroatom to a $-(CH_2)_p-$ chain, alkylene or alkenylene as defined above.

The term "polyhaloalkyl" as used herein refers to an "alkyl" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as CF_3CH_2 , CF_3 or $\text{CF}_3\text{CF}_2\text{CH}_2$.



The term "polyhaloalkyloxy" as used herein refers to an "alkoxy" or "alkyloxy" group as defined above which includes from 2 to 9, preferably from 2 to 5, halo substituents, such as F or Cl, preferably F, such as CF₃CH₂O, CF₃O or CF₃CF₂CH₂O.

5 of R^4 include the following groups:





10 wherein R^a, R^b and R^c are H, alkyl, aryl or arylalkyl;
however, R^aO cannot be HO.

Examples of such prodrug esters R⁴ include



Other examples of suitable prodrug esters R⁴ include



25 is H, alkyl, halogen or alkoxy, R^e is alkyl, aryl, arylalkyl or alkoxy, and n₁ is 0, 1 or 2.

Where the compounds of structure I are in acid form they may form a pharmaceutically acceptable salt

such as alkali metal salts such as lithium, sodium or potassium, alkaline earth metal salts such as calcium or magnesium as well as zinc or aluminum and other cations such as ammonium, choline, diethanolamine, lysine (D or L), ethylenediamine, t-butylamine, t-octylamine, tris-(hydroxymethyl)aminomethane (TRIS), N-methyl glucosamine (NMG), triethanolamine and dehydroabietylamine.

All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one or the R substituents. Consequently, compounds of formula I can exist in enantiomeric or diastereomeric forms or in mixtures thereof. The processes for preparation can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric or enantiomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional crystallization.

Where desired, the compounds of structure I may be used in combination with one or more hypolipidemic agents or lipid-lowering agents and/or one or more other types of therapeutic agents including antidiabetic agents, anti-obesity agents, antihypertensive agents, platelet aggregation inhibitors, and/or anti-osteoporosis agents, which may be administered orally in the same dosage form, in a separate oral dosage form or by injection.

The hypolipidemic agent or lipid-lowering agent which may be optionally employed in combination with the compounds of formula I of the invention may include 1,2,3 or more MTP inhibitors, HMG CoA reductase inhibitors, squalene synthetase inhibitors, fibric acid derivatives, ACAT inhibitors, lipoxxygenase inhibitors, cholesterol absorption inhibitors, ileal Na⁺/bile acid cotransporter inhibitors, upregulators of LDL receptor activity, bile


acid sequestrants, and/or nicotinic acid and derivatives thereof.

MTP inhibitors employed herein include MTP inhibitors disclosed in U.S. Patent No. 5,595,872, U.S. Patent No. 5,739,135, U.S. Patent No. 5,712,279, U.S. Patent No. 5,760,246, U.S. Patent No. 5,827,875, U.S. Patent No. 5,885,983 and U.S. Application Serial No. 09/175,180 filed October 20, 1998, now U.S. Patent No. 5,962,440. Preferred are each of the preferred MTP inhibitors disclosed in each of the above patents and applications.

All of the above U.S. Patents and applications are incorporated herein by reference.

Most preferred MTP inhibitors to be employed in accordance with the present invention include preferred MTP inhibitors as set out in U.S. Patent Nos. 5,739,135 and 5,712,279, and U.S. Patent No. 5,760,246.

The most preferred MTP inhibitor is 9-[4-[4-[[2-(2,2,2-Trifluoroethoxy)benzoyl]amino]-1-piperidinyl]butyl]-N-(2,2,2-trifluoroethyl)-9H-fluorene-9-carboxamide



The hypolipidemic agent may be an HMG CoA reductase inhibitor which includes, but is not limited to, mevastatin and related compounds as disclosed in U.S. Patent No. 3,983,140, lovastatin (mevinolin) and related compounds as disclosed in U.S. Patent No. 4,231,938, pravastatin and related compounds such as disclosed in U.S. Patent No. 4,346,227, simvastatin and related compounds as disclosed in U.S. Patent Nos. 4,448,784 and

The squalene synthetase inhibitors suitable for
35 use herein include, but are not limited to, α -phosphono-
sulfonates disclosed in U.S. Patent No. 5,712,396, those
disclosed by Biller et al, J. Med. Chem., 1988, Vol. 31,

No. 10, pp 1869-1871, including isoprenoid (phosphinyl-methyl)phosphonates as well as other known squalene synthetase inhibitors, for example, as disclosed in U.S. Patent No. 4,871,721 and 4,924,024 and in Biller, S.A., Neuenschwander, K., Ponpipom, M.M., and Poulter, C.D., Current Pharmaceutical Design, 2, 1-40 (1996).

In addition, other squalene synthetase inhibitors suitable for use herein include the terpenoid pyrophosphates disclosed by P. Ortiz de Montellano et al, J. Med. Chem., 1977, 20, 243-249, the farnesyl diphosphate analog A and presqualene pyrophosphate (PSQ-PP) analogs as disclosed by Corey and Volante, J. Am. Chem. Soc., 1976, 98, 1291-1293, phosphinylphosphonates reported by McClard, R.W. et al, J.A.C.S., 1987, 109, 5544 and cyclopropanes reported by Capson, T.L., PhD dissertation, June, 1987, Dept. Med. Chem. U of Utah, Abstract, Table of Contents, pp 16, 17, 40-43, 48-51, Summary.

Other hypolipidemic agents suitable for use herein include, but are not limited to, fibric acid derivatives, such as fenofibrate, gemfibrozil, clofibrate, bezafibrate, ciprofibrate, clinofibrate and the like, probucol, and related compounds as disclosed in U.S. Patent No. 3,674,836, probucol and gemfibrozil being preferred, bile acid sequestrants such as cholestyramine, colestipol and DEAE-Sephadex (Secholex®, Policexide®) and cholestagel (Sankyo/Geltex), as well as lipostabil (Rhone-Poulenc), Eisai E-5050 (an N-substituted ethanolamine derivative), imanixil (HOE-402), tetrahydrolipstatin (THL), istigmastanylphosphorylcholine (SPC, Roche), aminocyclodextrin (Tanabe Seiyoku), Ajinomoto AJ-814 (azulene derivative), melinamide (Sumitomo), Sandoz 58-035, American Cyanamid CL-277,082 and CL-283,546 (disubstituted urea derivatives), nicotinic acid (niacin), acipimox, acifran, neomycin, p-aminosalicylic acid, aspirin, poly(diallylmethylamine) derivatives such as disclosed in

U.S. Patent No. 4,759,923, quaternary amine poly(diallyldimethylammonium chloride) and ionenes such as disclosed in U.S. Patent No. 4,027,009, and other known serum cholesterol lowering agents.

- 5 The hypolipidemic agent may be an ACAT inhibitor such as disclosed in, Drugs of the Future 24, 9-15 (1999), (Avasimibe); "The ACAT inhibitor, Cl-1011 is effective in the prevention and regression of aortic fatty streak area in hamsters", Nicolosi et al, 10 Atherosclerosis (Shannon, Irel). (1998), 137(1), 77-85; "The pharmacological profile of FCE 27677: a novel ACAT inhibitor with potent hypolipidemic activity mediated by selective suppression of the hepatic secretion of ApoB100-containing lipoprotein", Ghiselli, Giancarlo, 15 Cardiovasc. Drug Rev. (1998), 16(1), 16-30; "RP 73163: a bioavailable alkylsulfinyl-diphenylimidazole ACAT inhibitor", Smith, C., et al, Bioorg. Med. Chem. Lett. (1996), 6(1), 47-50; "ACAT inhibitors: physiologic mechanisms for hypolipidemic and anti-atherosclerotic 20 activities in experimental animals", Krause et al, Editor(s): Ruffolo, Robert R., Jr.; Hollinger, Manfred A., Inflammation: Mediators Pathways (1995), 173-98, Publisher: CRC, Boca Raton, Fla.; "ACAT inhibitors: potential anti-atherosclerotic agents", Sliskovic et al, 25 Curr. Med. Chem. (1994), 1(3), 204-25; "Inhibitors of acyl-CoA:cholesterol O-acyl transferase (ACAT) as hypocholesterolemic agents. 6. The first water-soluble ACAT inhibitor with lipid-regulating activity. Inhibitors of acyl-CoA:cholesterol acyltransferase (ACAT). 7. 30 Development of a series of substituted N-phenyl-N'-[(1-phenylcyclopentyl)methyl]ureas with enhanced hypocholesterolemic activity", Stout et al, Chemtracts: Org. Chem. (1995), 8(6), 359-62, or TS-962 (Taisho Pharmaceutical Co. Ltd).
- 35 The hypolipidemic agent may be an upregulator of LD2 receptor activity such as MD-700 (Taisho Pharmaceutical Co. Ltd) and LY295427 (Eli Lilly).

The hypolipidemic agent may be a cholesterol absorption inhibitor preferably Schering-Plough's SCH48461 as well as those disclosed in Atherosclerosis 115, 45-63 (1995) and J. Med. Chem. 41, 973 (1998).

5 The hypolipidemic agent may be an ileal Na⁺/bile acid cotransporter inhibitor such as disclosed in Drugs of the Future, 24, 425-430 (1999).

10 The lipid-modulating agent may be a cholesteryl ester transfer protein (CETP) inhibitor such as Pfizer's CP 529,414 (WO/0038722 and EP 818448) and Pharmacia's SC-744 and SC-795.

15 The ATP citrate lyase inhibitor which may be employed in the combination of the invention may include, for example, those disclosed in U.S. Patent No. 5,447,954.

 Preferred hypolipidemic agents are pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin, cerivastatin, itavastatin and visastatin and ZD-4522.

20 The above-mentioned U.S. patents are incorporated herein by reference. The amounts and dosages employed will be as indicated in the Physician's Desk Reference and/or in the patents set out above.

25 The compounds of formula I of the invention will be employed in a weight ratio to the hypolipidemic agent (were present), within the range from about 500:1 to about 1:500, preferably from about 100:1 to about 1:100.

30 The dose administered must be carefully adjusted according to age, weight and condition of the patient, as well as the route of administration, dosage form and regimen and the desired result.

 The dosages and formulations for the hypolipidemic agent will be as disclosed in the various patents and applications discussed above.

35 The dosages and formulations for the other hypolipidemic agent to be employed, where applicable, will be as set out in the latest edition of the Physicians' Desk Reference.

For oral administration, a satisfactory result may be obtained employing the MTP inhibitor in an amount within the range of from about 0.01 mg to about 500 mg and preferably from about 0.1 mg to about 100 mg, one to
5 four times daily.

A preferred oral dosage form, such as tablets or capsules, will contain the MTP inhibitor in an amount of from about 1 to about 500 mg, preferably from about 2 to about 400 mg, and more preferably from about 5 to about
10 250 mg, one to four times daily.

For oral administration, a satisfactory result may be obtained employing an HMG CoA reductase inhibitor, for example, pravastatin, lovastatin, simvastatin, atorvastatin, fluvastatin or cerivastatin in dosages
15 employed as indicated in the Physician's Desk Reference, such as in an amount within the range of from about 1 to 2000 mg, and preferably from about 4 to about 200 mg.

The squalene synthetase inhibitor may be employed in dosages in an amount within the range of from about 10
20 mg to about 2000 mg and preferably from about 25 mg to about 200 mg.

A preferred oral dosage form, such as tablets or capsules, will contain the HMG CoA reductase inhibitor in an amount from about 0.1 to about 100 mg, preferably from
25 about 0.5 to about 80 mg, and more preferably from about 1 to about 40 mg.

A preferred oral dosage form, such as tablets or capsules will contain the squalene synthetase inhibitor in an amount of from about 10 to about 500 mg, preferably
30 from about 25 to about 200 mg.

The hypolipidemic agent may also be a lipoxxygenase inhibitor including a 15-lipoxxygenase (15-LO) inhibitor such as benzimidazole derivatives as disclosed in WO 97/12615, 15-LO inhibitors as disclosed in WO 97/12613,
35 isothiazolones as disclosed in WO 96/38144, and 15-LO inhibitors as disclosed by Sendobry et al "Attenuation of diet-induced atherosclerosis in rabbits with a highly

5

10

15

cerivastatin as well as niacin and/or cholestagel.

20

3.

compounds of structure I will be employed in a weight

The other antidiabetic agent may also preferably be a sulfonyl urea such as glyburide (also known as glibenclamide), glimepiride (disclosed in U.S. Patent No. 4,379,785), glipizide, gliclazide or chlorpropamide, other known sulfonylureas or other antihyperglycemic agents which act on the ATP-dependent channel of the β -cells, with glyburide and glipizide being preferred, which may be administered in the same or in separate oral dosage forms.

The oral antidiabetic agent may also be a glucosidase inhibitor such as acarbose (disclosed in U.S. Patent No. 4,904,769) or miglitol (disclosed in U.S. Patent No. 4,639,436), which may be administered in the same or in a separate oral dosage forms.

The compounds of structure I may be employed in combination with a PPAR γ agonist such as a thiazolidinedione oral anti-diabetic agent or other insulin sensitizers (which has an insulin sensitivity effect in NIDDM patients) such as troglitazone (Warner-Lambert's Rezulin[®], disclosed in U.S. Patent No. 4,572,912), rosiglitazone (SKB), pioglitazone (Takeda), Mitsubishi's MCC-555 (disclosed in U.S. Patent No. 5,594,016), Glaxo-Welcome's GL-262570, englitazone (CP-68722, Pfizer) or darglitazone (CP-86325, Pfizer, isaglitazone (MIT/J&J), JTT-501 (JPNT/P&U), L-895645 (Merck), R-119702 (Sankyo/WL), NN-2344 (Dr. Reddy/NN), or

YM-440 (Yamanouchi), preferably rosiglitazone and pioglitazone.

5 The compounds of structure I will be employed in a weight ratio to the thiazolidinedione in an amount within the range from about 0.01:1 to about 100:1, preferably from about 0.05 to about 10:1.

10 The sulfonyl urea and thiazolidinedione in amounts of less than about 150 mg oral antidiabetic agent may be incorporated in a single tablet with the compounds of structure I.

15 The compounds of structure I may also be employed in combination with a antihyperglycemic agent such as insulin or with glucagon-like peptide-1 (GLP-1) such as GLP-1(1-36) amide, GLP-1(7-36) amide, GLP-1(7-37) (as disclosed in U.S. Patent No. 5,614,492 to Habener, the disclosure of which is incorporated herein by reference), as well as AC2993 (Amylin) and LY-315902 (Lilly), which may be administered via injection, intranasal, inhalation or by transdermal or buccal devices.

20 Where present, metformin, the sulfonyl ureas, such as glyburide, glimepiride, glipyrider, glipizide, chlorpropamide and gliclazide and the glucosidase inhibitors acarbose or miglitol or insulin (injectable, pulmonary, buccal, or oral) may be employed in
25 formulations as described above and in amounts and dosing as indicated in the Physician's Desk Reference (PDR).

30 Where present, metformin or salt thereof may be employed in amounts within the range from about 500 to about 2000 mg per day which may be administered in single or divided doses one to four times daily.

Where present, the thiazolidinedione anti-diabetic agent may be employed in amounts within the range from about 0.01 to about 2000 mg/day which may be administered in single or divided doses one to four times per day.

35 Where present insulin may be employed in formulations, amounts and dosing as indicated by the Physician's Desk Reference.

5 incorporated herein by reference.

15 The antidiabetic agent may be an SGLT2 inhibitor
such as disclosed in U.S. provisional application No.
60/158,773, filed October 12, 1999 (attorney file LA49),
employing dosages as set out therein. Preferred are the
compounds designated as preferred in the above
20 application.

The antidiabetic agent may be a DP4 inhibitor such as disclosed in Provisional Application 60/188,555 filed March 10, 2000 (attorney file LA50), WO99/38501, WO99/46272, WO99/67279 (PROBIODRUG), WO99/67278 (PROBIODRUG), WO99/61431 (PROBIODRUG), NVP-DPP728A (1-[[[2-[(5-cyanopyridin-2-yl)amino]ethyl]amino]acetyl]-2-cyano-(S)-pyrrolidine) (Novartis) (preferred) as disclosed by Hughes et al, Biochemistry, 38(36), 11597-11603, 1999, TSL-225 (tryptophyl-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid (disclosed by Yamada et

al, Bioorg. & Med. Chem. Lett. 8 (1998) 1537-1540, 2-cyanopyrrolidides and 4-cyanopyrrolidides as disclosed by Ashworth et al, Bioorg. & Med. Chem. Lett., Vol. 6, No. 22, pp 1163-1166 and 2745-2748 (1996) employing dosages
5 as set out in the above references.

The meglitinide which may optionally be employed in combination with the compound of formula I of the invention may be repaglinide, nateglinide (Novartis) or KAD1229 (PF/Kissei), with repaglinide being preferred.

10 The compound of formula I will be employed in a weight ratio to the meglitinide, PPAR γ agonist, PPAR α/γ dual agonist, $\alpha P2$ inhibitor, DP4 inhibitor or SGLT2 inhibitor within the range from about 0.01:1 to about 100:1, preferably from about 0.05 to about 10:1.

15 The other type of therapeutic agent which may be optionally employed with a compound of formula I may be 1, 2, 3 or more of an anti-obesity agent including a beta 3 adrenergic agonist, a lipase inhibitor, a serotonin (and dopamine) reuptake inhibitor, an $\alpha P2$ inhibitor, a
20 thyroid receptor agonist and/or an anorectic agent.

The beta 3 adrenergic agonist which may be optionally employed in combination with a compound of formula I may be AJ9677 (Takeda/Dainippon), L750355 (Merck), or CP331648 (Pfizer) or other known beta 3
25 agonists as disclosed in U.S. Patent Nos. 5,541,204, 5,770,615, 5,491,134, 5,776,983 and 5,488,064, with AJ9677, L750,355 and CP331648 being preferred.

The lipase inhibitor which may be optionally employed in combination with a compound of formula I may
30 be orlistat or ATL-962 (Alizyme), with orlistat being preferred.

The serotonin (and dopamine) reuptake inhibitor which may be optionally employed in combination with a compound of formula I may be sibutramine, topiramate
35 (Johnson & Johnson) or axokine (Regeneron), with sibutramine and topiramate being preferred.

The thyroid receptor agonist which may be optionally employed in combination with a compound of formula I may be a thyroid receptor ligand as disclosed in WO97/21993 (U. Cal SF), WO99/00353 (KaroBio),
5 GB98/284425 (KaroBio), and U.S. Provisional Application 60/183,223 filed February 17, 2000, with compounds of the KaroBio applications and the above U.S. provisional application being preferred.

The anorectic agent which may be optionally
10 employed in combination with a compound of formula I may be dexamphetamine, phentermine, phenylpropanolamine or mazindol, with dexamphetamine being preferred.

The various anti-obesity agents described above may be employed in the same dosage form with the compound of
15 formula I or in different dosage forms, in dosages and regimens as generally known in the art or in the PDR.

The antihypertensive agents which may be employed in combination with the compound of formula I of the invention include ACE inhibitors, angiotensin II receptor
20 antagonists, NEP/ACE inhibitors, as well as calcium channel blockers, β -adrenergic blockers and other types of antihypertensive agents including diuretics.

The angiotensin converting enzyme inhibitor which may be employed herein includes those containing a
25 mercapto (-S-) moiety such as substituted proline derivatives, such as any of those disclosed in U.S. Pat. No. 4,046,889 to Ondetti et al mentioned above, with captopril, that is, 1-[(2S)-3-mercapto-2-methylpropionyl]-L-proline, being preferred, and
30 mercaptoacyl derivatives of substituted prolines such as any of those disclosed in U.S. Pat. No. 4,316,906 with zofenopril being preferred.

Other examples of mercapto containing ACE inhibitors that may be employed herein include rentiapril
35 (fentiapril, Santen) disclosed in Clin. Exp. Pharmacol. Physiol. 10:131 (1983); as well as pivopril and YS980.

Other examples of angiotensin converting enzyme inhibitors which may be employed herein include any of those disclosed in U.S. Pat. No. 4,374,829 mentioned above, with N-(1-ethoxycarbonyl-3-phenylpropyl)-L-alanyl-L-proline, that is, enalapril, being preferred, any of the phosphonate substituted amino or imino acids or salts disclosed in U.S. Pat. No. 4,452,790 with (S)-1-[6-amino-2-[[hydroxy-(4-phenylbutyl)phosphinyl]oxy]-1-oxohexyl]-L-proline or (ceronapril) being preferred, phosphinylalkanoyl prolines disclosed in U.S. Pat. No. 4,168,267 mentioned above with fosinopril being preferred, any of the phosphinylalkanoyl substituted prolines disclosed in U.S. Pat. No. 4,337,201, and the phosphonamides disclosed in U.S. Pat. No. 4,432,971 discussed above.

Other examples of ACE inhibitors that may be employed herein include Beecham's BRL 36,378 as disclosed in European Patent Application Nos. 80822 and 60668; Chugai's MC-838 disclosed in C.A. 102:72588v and Jap. J. Pharmacol. 40:373 (1986); Ciba-Geigy's CGS 14824 (3-([1-ethoxycarbonyl-3-phenyl-(1S)-propyl]amino)-2,3,4,5-tetrahydro-2-oxo-1-(3S)-benzazepine-1 acetic acid HCl) disclosed in U.K. Patent No. 2103614 and CGS 16,617 (3(S)-[[(1S)-5-amino-1-carboxypentyl]amino]-2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-ethanoic acid) disclosed in U.S. Pat. No. 4,473,575; cetapril (alacepril, Dainippon) disclosed in Eur. Therap. Res. 39:671 (1986); 40:543 (1986); ramipril (Hoechst) disclosed in Euro. Patent No. 79-022 and Curr. Ther. Res. 40:74 (1986); Ru 44570 (Hoechst) disclosed in Arzneimittelforschung 34:1254 (1985), cilazapril (Hoffman-LaRoche) disclosed in J. Cardiovasc. Pharmacol. 9:39 (1987); R 31-2201 (Hoffman-LaRoche) disclosed in FEBS Lett. 165:201 (1984); lisinopril (Merck), indalaprill (delapril) disclosed in U.S. Pat. No. 4,385,051; indolapril (Schering) disclosed in J. Cardiovasc. Pharmacol. 5:643, 655 (1983), spirapril (Schering)

5

10

15

30

35

5 more preferably from about 10 to about 150 mg.

10 mg/kg to about 1 mg/kg.

conventional carriers.

15 It will be appreciated that preferred dosages of ACE inhibitor and AII antagonist as well as other antihypertensives disclosed herein will be as set out in the latest edition of the Physician's Desk Reference (PDR).

20 Other examples of preferred antihypertensive agents
suitable for use herein include omapatrilat (Vanlev®)
amlodipine besylate (Norvasc®), prazosin HCl
(Minipress®), verapamil, nifedipine, nadolol, diltiazem,
felodipine, nisoldipine, isradipine, nicardipine,
25 atenolol, carvedilol, sotalol, terazosin, doxazosin,
propranolol, and clonidine HCl (Catapres®).

torasemide, furosemide, spironolactone, and indapamide.

30 Antiplatelet agents which may be employed in combination with compounds of formula I of the invention include aspirin, clopidogrel, ticlopidine, dipyridamole, abciximab, tirofiban, eptifibatide, anagrelide, and ifetroban, with clopidogrel and aspirin being preferred.

35 The antiplatelet drugs may be employed in amounts
as indicated in the PDR. Ifetroban may be employed in
amounts as set out in U.S. Patent No. 5,100,889.

Antiosteoporosis agents suitable for use herein in combination with the compounds of formula I of the invention include parathyroid hormone or bisphosphonates, such as MK-217 (alendronate) (Fosamax®). Dosages
5 employed will be as set out in the PDR.

In carrying out the method of the invention, a pharmaceutical composition will be employed containing the compounds of structure I, with or without another therapeutic agent, in association with a pharmaceutical
10 vehicle or diluent. The pharmaceutical composition can be formulated employing conventional solid or liquid vehicles or diluents and pharmaceutical additives of a type appropriate to the mode of desired administration. The compounds can be administered to mammalian species
15 including humans, monkeys, dogs, etc. by an oral route, for example, in the form of tablets, capsules, granules or powders, or they can be administered by a parenteral route in the form of injectable preparations. The dose for adults is preferably between 50 and 2,000 mg per day,
20 which can be administered in a single dose or in the form of individual doses from 1-4 times per day.

A typical capsule for oral administration contains compounds of structure I (250 mg), lactose (75 mg) and magnesium stearate (15 mg). The mixture is passed
25 through a 60 mesh sieve and packed into a No. 1 gelatin capsule.

A typical injectable preparation is produced by aseptically placing 250 mg of compounds of structure I into a vial, aseptically freeze-drying and sealing. For
30 use, the contents of the vial are mixed with 2 mL of physiological saline, to produce an injectable preparation.

The following Examples represent preferred
35 embodiments of the invention.

-


40

CC1=C(C(=N1C2=CC=CC=C2)OCCOC3=CC=C(C=C3)CN(Cc4ccccc4)C(=O)O)CCC1=C(C(=N1C2=CC=CC=C2)CCOC3=CC=C(C=C3)C=O)OC4=CC=CC=C4

To a 0°C solution of 4-hydroxybenzaldehyde (1.70 g, 12.3 mmol), 5-phenyl-2-methyl-oxazole-4-ethanol (Maybridge; 2.50 g, 14.0 mmol) and Ph₃P (4.20 g, 16.0 mmol) in dry THF (30 mL) was added dropwise DEAD (3.20 g, 15.0 mmol). The solution was stirred at 0°C for 0.5 h, then was allowed to warm to RT and stirred overnight. The orange-red solution was concentrated *in vacuo* and the residue was chromatographed (stepwise gradient from 5:1 to 5:2 hex:EtOAc) to give Part A compound (2.47 g, 65%) as a clear, slightly yellow viscous oil.

Al. Alternative Procedure for Preparing Part A Aldehyde


(1)



To a -5°C solution of 5-phenyl-2-methyl-oxazole-4-ethanol (20.00 g, 0.098 mol) in CH₂Cl₂ (100 mL) was added methanesulfonyl chloride (12.40 g, 0.108 mol) in one portion (exothermic reaction). After recooling to -5°C, Et₃N (11.1 g, 0.110 mol) was added slowly over 30 min (internal temperature <3°C). The reaction was allowed to warm to RT and stirred for 1 h (reaction monitored by analytical HPLC), at which point starting material had been consumed. The reaction was washed with aqueous HCl (2 x 50 mL of a 3N solution). The combined aqueous layers were extracted with CH₂Cl₂ (50 mL). The combined organic extracts were successively washed with satd. aqueous NaHCO₃ and brine (50 mL each), dried (Na₂SO₄), and concentrated to ~30 mL volume. Methyl tert-butyl ether (120 mL) was added and the mixture was stirred; a white solid was formed. The mixture was cooled to -20°C for

complete crystallization. The product was filtered and vacuum-dried to give the product mesylate (23.3 g, 85%) as a white solid. The mother liquor was concentrated in vacuo and recrystallized from methyl tert butyl ether/heptane to give a second crop of product mesylate (3.3 g, 12%; total yield = 97%).

(2)




10

A mixture of the above mesylate (13.6 g, 0.048 mol), 4-hydroxybenzaldehyde (7.09 g, 0.058 mol) and K_2CO_3 (9.95 g, 0.072 mol) in DMF (110 mL) was heated at 100°C for 2 h (reaction complete by analytical HPLC). The mixture was allowed to cool to RT and then poured into ice-water (400 mL) and stirred for 30 min. The solid product was filtered and washed with cold water (3 x 25 mL) and dried in vacuo at 50°-60°C overnight. The crude product was crystallized from MTBE-Hexane to give (12.2 g, 82%; 2 crops) the aldehyde (Part A1 compound) as a white solid.

20

B.



25

To a solution of N-benzyl glycine ethyl ester (43 mg; 0.22 mmol) and Part A1 compound (52 mg; 0.17 mmol) in DCE (10 mL) was added $NaBH(OAc)_3$ (56 mg; 0.26 mmol). The reaction mixture was stirred vigorously overnight for 12 hours. Saturated aqueous $NaHCO_3$ (10 mL) was added, and the mixture was extracted with EtOAc (3x 10 mL). The

30


5

CC1=C(C(=N1C2=CC=CC=C2)OCCOC3=CC=C(C=C3)CN(Cc4ccccc4)C(=O)O)C5=CC=CC=C5


To a solution of Part B compound (45 mg) in MeOH (2 mL) was added aqueous NaOH (3 mL of a 1M solution). The solution was stirred overnight for 14 h and then acidified to pH 5 with excess aqueous HCl (1M solution).

15

Example 2




- 90 -




To a solution of Example 1 Part A compound (147 mg;
5 0.479 mmol) and glycine ethyl ester hydrochloride (73 mg;
0.52 mmol) in DCE (2 mL) was added Et₃N and NaBH(OAc)₃
(156 mg; 0.74 mmol) and the reaction was stirred
overnight at RT. Flash chromatography (stepwise gradient
from 7:3 to 2:3 hex: EtOAc) gave 35 mg (21%) of the
10 dibenzyl glycine ester (Example 2 Part A compound). In
addition, 127 mg (67%) of the monobenzyl glycine ester
(Example 3 Part A compound) was obtained.


B.



5 A solution of Example 1 Part A compound (35 mg;
0.051 mmol) in MeOH (2 mL) and aqueous NaOH (3 mL of a 1M
solution) was heated under reflux for 12 h. The solution
was adjusted to pH 5 with aqueous 1M HCl and aqueous 1 M
NaOH, then extracted with EtOAc (3x). The combined
10 organic extracts were washed with brine, dried (Na₂SO₄),
and concentrated in vacuo to give title compound (13 mg)
as a colorless solid. [M + H]⁺ = 658.2


Example 3

A.




20 To a solution of Example 1 Part A compound (147 mg;
0.479 mmol) and glycine ethyl ester hydrochloride (73 mg;
mmol) in DCE was added Et₃N and NaBH(OAc)₃ (156 mg; 0.74
mmol). Flash chromatography (stepwise gradient from 7:3
25 to 2:3 hex: EtOAc) gave 127 mg (67%) of the title
compound. In addition, 35 mg (21%) of the bis-benzyl
glycine ester (Example 2 Part A compound) was obtained as
a byproduct.

B.




5 A solution of Part A compound (72 mg; 0.18 mmol) in aqueous NaOH (2 mL of a 1M solution) and MeOH (2 mL) was refluxed for 3 h. The reaction was adjusted to pH 5 with aqueous 1M HCl, and solids were filtered off. The filtrate was extracted with EtOAc (3x). The combined
 10 organic extracts were washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to give a colorless solid, which was purified by preparative HPLC (utilizing a YMC S5 ODS 20 mm x 100 mm column with a continuous gradient from 70% A:30 %B to 100% B for 10 min at a flow rate of
 15 20 mL/min, where A = 90:10:0.1 H₂O:MeOH:TFA and where B = 90:10:0.1 MeOH:H₂O:TFA) to give title compound (10 mg; 15%) as a colorless solid. [M + H]⁺ = 367.2

Example 4

20


A solution of the amino t-butyl ester (0.040 g, 0.095 mmol), (prepared as described for Example 7 Part C, except that the aldehyde used in the reductive amination was Example 1 Part A instead of Example 7 Part A)
 25



and propargyl bromide (0.014 g, 0.120 mmol) and DBU (0.5 mL; 2.96 mmol) in DCE (1 mL) was stirred at 0°C for 5 h.

TLC showed that the reaction was complete at this point. EtOAc (10 mL) was added and the organic phase was washed with H₂O and concentrated *in vacuo*. The residual oil was dissolved in CH₂Cl₂/TFA (1:1, 1 mL) and stirred at RT for 5 h, then concentrated *in vacuo*. The residue was purified by preparative HPLC (YMC S5 ODS 30 mm x 250 mm reverse phase column; flow rate = 25 mL/min; 30 min continuous gradient from 70:30 A:B to 100% B; where A = 90:10:0.1 H₂O:MeOH:TFA and where B = 90:10:0.1 MeOH:H₂O:TFA) to give the title compound (34 mg, 92%) as an oil. LC/MS (electrospray) gave the correct [M + H]⁺ = 405.2 for the title compound.


Example 5



15

A solution of 2-chlorobenzoxazole (20 mg; 0.131 mmol), the secondary amine-methyl ester (52 mg; 0.146 mmol)

20




(prepared as described in Example 3 Part A except glycine ethyl ester HCl was replaced by glycine methyl ester HCl and the Example 7 Part A aldehyde was employed), and excess Et₃N (0.5 mL) in THF (2.0 mL) was heated to 100°C in a sealed tube and the reaction was monitored by LC/MS. After 4 days, starting amine had been consumed. The reaction was cooled to RT and aqueous LiOH (0.50 mL of a 1 M solution) was added to the solution. The solution was stirred at RT for 5 h, after which the hydrolysis was

30

10

CC1=C(C(=O)O)C(=O)N1Cc2ccc(OCCc3c(C)c(C(=O)O)nc3C4=CC=CC=C4)cc2

15




20

[illegible]


25

A.



5 To a 0°C solution of 3-hydroxybenzaldehyde (3.00 g; 24.6 mmol), 2-phenyl-5-methyl-oxazole-4-ethanol (5.00 g; 24.6 mmol) and Ph₃P (7.10 g; 27.1 mmol) in dry THF (75 mL) was added dropwise DEAD (4.27 mL; 27.1 mmol) over 10 min. The brown-orange solution was allowed to warm to RT and stirred at RT for 24 h. The solution was concentrated *in vacuo* and chromatographed (SiO₂; stepwise gradient: 100% hex to hex:EtOAc 3:1) to give Part A compound as a pale yellow viscous oil (4.01 g; 53%).

15 A.1. Alternative Procedure for Preparing Part A Aldehyde




20 To a solution of 3-hydroxybenzaldehyde (9.1 g; 0.074 mmol) in CH₃CN (206 mL) was added K₂CO₃ (10.3 g). The mixture was heated to 90°C in an oil bath and stirred for 18 h at 90°C (the reaction was complete at this point by analytical HPLC). The reaction was cooled to RT, then diluted with EtOAc (500 mL), washed with H₂O, aqueous NaOH (2 x 100 mL of a 1 M solution) and brine. The organic phase was dried (MgSO₄) and concentrated *in vacuo*. The residual oil was chromatographed (SiO₂; hex:EtOAc from 9:1 to 4:1) to give the Part A aldehyde (12.7 g; 67%) as a viscous, clear, pale yellow oil.

25


30

B.



5 A solution of the Part A1 compound (4.00 g; 13.0 mmol), glycine tert-butyl ester hydrochloride (2.40 g; 14.3 mmol) and Et₃N (2.18 mL; 15.7 mmol) in MeOH (30 mL) was stirred at RT for 6 h and then cooled to 0°C. A solution of NaBH₄ (594 mg; 15.7 mmol) in MeOH (10 mL) was
10 added portionwise at 0°C to the solution of crude imine over ~15 min. The solution was stirred at 0°C for 3 h, then at RT for 3 h, then concentrated *in vacuo* without heating to removed MeOH. The residue was partitioned between saturated aqueous NaCl and EtOAc (50 mL each).
15 The aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic extracts were dried (Na₂SO₄) and concentrated *in vacuo* to give a yellow oil, which was chromatographed on SiO₂ (stepwise gradient; hex:EtOAc from 4:1 to 2:3) to give Part B compound as a pale
20 viscous yellow oil (4.82 g; 88%).

C.




25

To a solution of Part B compound (0.400 g; 0.95 mmol) and 4-phenoxybenzaldehyde (0.216 g; 1.09 mmol) in DCE (5 mL) was added NaBH(OAc)₃ (0.300 g; 1.42 mmol), followed by HOAc (25 µL). The reaction was stirred at RT
30 for 24 h. 10% unreacted starting amine was still present by analytical HPLC. Additional aldehyde (30 mg) and


NaBH(OAc)₃ (60 mg) were added and the reaction was stirred at RT for a further 18 h, after which reaction was complete. The solution was partitioned between aqueous NaHCO₃ (50 mL of a 10% solution) and EtOAc (50 mL). The aqueous layer was extracted with EtOAc (2 x 25 mL). The combined organic extracts were washed with aqueous NaHCO₃ (2 x 15 mL of a 10% solution), dried (Na₂SO₄) and concentrated *in vacuo* to give Part C compound (521 mg crude material) as a clear, colorless oil.

D.




Part C compound was dissolved in CHCl₃ (2 mL) and TFA (1.5 mL) and the solution was stirred at RT for 24 h. The solution was concentrated *in vacuo* and the residue was purified by preparative HPLC (YMC S5 ODS 20 x 250 mm column; continuous gradient from 40:60 solvent A:B to 100% solvent B; where solvent A = 90:10:0.1 H₂O:MeOH:TFA; solvent B = 90:10:0.1 MeOH:H₂O:TFA). The purified product was lyophilized from MeOH/H₂O to give the title amino acid (312 mg; 48% over 2 steps) as its TFA salt (off-white lyophilate). [M+H]⁺ (electrospray) = 549.3

Example 8




A mixture of the amino-ester (39 mg; 0.092 mmol),




(prepared as described in Example 4),

2-naphthaldehyde (29 mg; 0.185 mmol), and $\text{NaBH}(\text{OAc})_3$ (100 mg; 0.472 mmol) in DCE (1.5 mL) was stirred at RT for 16 h. TFA (1.0 mL) was then added to the mixture, which was stirred at RT for a further 12 h. Volatiles were removed *in vacuo*. The resulting residue was diluted with MeOH (1.5 mL), filtered, and purified by preparative HPLC (YMC S5 ODS 30mm x 250 mm column; continuous 30 min gradient @ 25 mL/min from 100% A to 100% B; solvent A = 90:10:0.1 $\text{H}_2\text{O}:\text{MeOH}:\text{TFA}$; B = 90:10:0.1 $\text{MeOH}:\text{H}_2\text{O}:\text{TFA}$) to give the desired title product (39 mg; 68%) as a clear, viscous oil. $[\text{M} + \text{H}]^+ = 507.3$


Example 9



A.




A solution of the amino acid tert-butyl ester (1.8 g, 4.27 mmol)



(prepared as described in Example 7 Part B), and TFA (20 mL) in CH_2Cl_2 (40 mL) was stirred at RT overnight. The solution was concentrated *in vacuo*, and

the residue was dissolved in CH_2Cl_2 and eluted through solid NaHCO_3 (to remove excess TFA) with excess CH_2Cl_2 . The combined filtrates were concentrated *in vacuo* to provide the desired amino acid Part A compound (1.48 g; 95%). $[\text{M} + \text{H}]^+ = 457.2$

B.



10

The title compound was prepared as part of a solution phase library run using the following exemplary procedure:

To a solution of the Part A amino acid compound (27 mg, 0.074 mmol; in 2 mL CH_2Cl_2) was added (4-chlorophenoxy)-3-benzaldehyde (86 mg; 0.37 mmol), $\text{NaBH}(\text{OAc})_3$ (79 mg, 0.37 mmol) and HOAc (0.1 mL). The reaction was stirred at RT for 15 h.

The product was purified via solid-phase extraction using a Varian SAX cartridge (3 g of sorbent in a 6 mL column, 0.3 meq/g) by the procedure outlined below:

- 1) The column was conditioned with MeOH (10 mL) and CH_2Cl_2 (20 mL)
- 2) The reaction mixture was loaded onto the SAX column
- 3) The column was rinsed with CH_2Cl_2 (10 mL)
- 4) The column was rinsed with 1% TFA in MeOH (3 mL)
- 5) The product was eluted with 1% TFA in MeOH (20 mL)


10

(procedure used with heterocyclic aldehydes)



20

A mixture of the amino acid (14 mg; 0.038 mmol),



25

5 68%) as a clear, viscous oil.

CC1=C(C(=N1C2=CC=CC=C2)OC3=CC=CC=C3)CCOC4=CC=CC=C4CN(CCC(=O)O)Cc5cc(oc5-c6ccc(Cl)cc6)O

An alternative purification procedure to preparative HPLC was used as follows:

15 Chemicals; 3 g of sorbent in a 6 mL column, 0.3 meq/g) by
the procedure outlined below:

- (5 mL)

2) The reaction mixture (diluted with 2 mL CH₂Cl₂) was loaded onto the SAX column

25

4) The product was eluted with 1% TFA in MeOH (20 mL)

30 product. This was dissolved in CH_2Cl_2 :MeOH (95:5) and loaded onto a silica gel cartridge (1.5 g SiO_2) and the product was eluted with CH_2Cl_2 :MeOH (95:5; 8 mL). The


Reverse Phase HPLC analysis (YMC S5 ODS 4.6 x 33 mm column, continuous gradient from 100% A to 100%B for 2 min at a flow rate of 5 mL/min [Solvent A=10% MeOH/90% H₂O/0.2% H₃PO₄; Solvent B = 90% MeOH/10% H₂O/0.2% H₃PO₄]) indicated that the product purity was 92%. In addition, LC/MS (electrospray) gave the correct molecular ion [(M+H)⁺ = 583] for title compound.

10

CCOC(=O)CNCCc1ccc(OCCc2c(C)c(O)c(c2N)C3=CC=CC=C3)cc1

20

B.



5

The title compound was prepared as part of a solution phase library run using the following exemplary procedure.

To a solution of Part A phenol compound (30 mg, 0.057 mmol) in CH_2Cl_2 (1 mL) was added 3-fluorophenyl boronic acid (12 mg; 0.086 mmol) and 4A molecular sieves (pre-dried at 400°C overnight) at RT. After stirring for 5 min, $\text{Cu}(\text{OAc})_2$ (1 eq), Et_3N (5 eq) and pyridine (5 eq) were added to the mixture. The vial was capped and air was allowed to pass into the reaction. The reaction was stirred at RT for 60 h and was complete by analytical HPLC and LC/MS. (For other reactions which were incomplete after this time, additional boronic acid (1.5 equivalent) was added in order to form additional desired product). The reaction mixture was filtered and concentrated *in vacuo*.

The product was purified via solid-phase extraction using a United Technology SCX column (2 g of sorbent in a 6 mL column) by the procedure outlined below.

25

1) The column was conditioned with MeOH (10 mL) and CH_2Cl_2 (10 mL)

2) The residue was dissolved in a minimal volume of CH_2Cl_2 and loaded onto the SCX column.


30

3) The cartridge was successively washed with CH_2Cl_2 (20 mL), $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (20% MeOH, 20 mL) and MeOH (20 mL)


4) The product was eluted with a solution of 0.5N NH_3 in MeOH.

5 The product-containing fractions were concentrated *in vacuo* to give the desired tert-butyl ester. (Some incomplete reactions required chromatography (on SiO_2) of the crude material to give esters of the requisite purity). The t-butyl ester was treated with a solution
10 of 30% TFA in CH_2Cl_2 overnight. Volatiles were removed and the residue was redissolved in CH_2Cl_2 (1 mL) and concentrated *in vacuo* on a Speed Vac to afford the desired title product (30 mg; 77%). Reverse phase HPLC analysis indicated that the product purity was 90%. In
15 addition LC/MS gave the correct molecular ion $[(\text{M}+\text{H})^+ = 567]$ for the desired title compound.


Example 12



A.



25 To a solution of the secondary amine-tert butyl ester (110 mg; 0.26 mmol)



(prepared as described in Example 7, Part B),

5

10



15

20

24

30

2) The residue was dissolved in a minimal volume of CH_2Cl_2 and loaded onto the SCX column.

5 4) The product was eluted with a solution of 0.5N NH_3 in
MeOH.

10

15


15

20 The t-butyl ester was treated with a solution of 1:1 TFA in CH₂Cl₂ overnight. Volatiles were removed and the residue was redissolved in CH₂Cl₂ (1 mL) and concentrated in vacuo on a Speed Vac to afford the desired title product (25 mg; 48%) as a slightly yellowish oil.

25

CC1=C(C(=O)N1Cc2ccccc2)C(=O)c3ccccc3

To a solution of 3-bromopyridine (32 mg; 0.2 mmol) in DME (1 mL) were successively added $(\text{Ph}_3\text{P})_4\text{Pd}$ (5 mg; 0.05 mol equiv) and the Example 12 Part A boronic acid (50 mg; 0.09 mmol)




The reaction mixture was filtered and the filtrate was chromatographed on a silica gel cartridge (2 mL; EtOAc). The product-containing fractions were concentrated *in vacuo* and the residue was chromatographed on another silica gel cartridge (2 mL; stepwise gradient of hexanes, hex:EtOAc 3:1 and EtOAc). The product-containing fractions were concentrated *in vacuo* and the residue was eluted through an SCX (2 g) cartridge (20 mL each of CH₂Cl₂ and MeOH; then product eluted with 2M ammonia in MeOH). The product-containing fractions were concentrated *in vacuo* to give the desired biaryl amine tert-butyl ester product. This was treated with a solution of CH₂Cl₂/TFA (7:3; 1 mL) overnight for 14 h. Volatiles were removed to give title compound (39 mg; 67%) as an oil. $[M + H]^+ = 534.3$











Examples 14 to 124












Following one of the above procedures, the following compounds of the invention were prepared:











Table 1



5

Example No.	R ³	[M+H] ⁺
14		457.3
15		471.3
16		485.3
17		617.2
18		549.3
19		533.3
20		557.3
21		617.3
22		562.7
23		579.3














Example No.	R ³	[M+H] ⁺
33		517.4
34		507.1
35		507.1
36		496.1
37		557.1
38		591.2
39		568.2
40		625.2
41		591.2
42		568.2
43		622.3














Example No.	R ³	[M+H] ⁺
44		601.2
45		557.2
46		519.2
47		675.2
48		519.2
4		600.3
50		564.2
51		545.3
52		625.2
53		563.3














CC1=C(C(=N1C2=CC=CC=C2)COC3=CC=CC=C3CN(CCC(=O)O)R3)C


- 114 -

Example No.	R ³	[M+H] ⁺
73		507.3
74		587.4
75		517.3
76		549.3
77		549.3
78		583.2
79		617.2
80		563.2
81		559.2
82		615.2
83		629.1
84		605.3
85		563.2


Example No.	R ³	[M+H] ⁺
86		596.2
87		549.3
88		635.3
89		639.2
90		583.2
91		563.2
92		635.3
93		583.2
94		617.2
95		617.1
96		567.2
97		555.1
98		595.3

Example No.	R ³	[M+H] ⁺
99		555.2
100		617.2
101		594.2
102		548.2
103		523.3
104		534.4
105		576.2
106		601.1
107		563.2
108		609.2
109		551.2
110		523.2
111		539.2

Example No.	R ³	[M+H] ⁺
112		579.3
113		594.4
114		563.3
115		583.2
116		579.3
117		583.2
118		594.3
119		594.3
120		537.3
121		537.3
122		535.2
123		535.2
124		496.1


Example 125

5 A.



A solution of Example 7 Part A aldehyde (60 mg; 0.20
10 mmol) and (S)- α -methyl benzylamine (30 mg; 0.24 mmol) in
MeOH (1 mL) was stirred at RT for 6 h. The solution was
cooled to 0°C and a pre-formed solution of NaBH₄ (9 mg;
0.24 mmol) in MeOH (0.5 mL) was added portionwise. The
reaction was stirred at RT overnight, then concentrated
15 in *vacuo* without heating. The residue was partitioned
between aqueous NaHCO₃ and EtOAc (5 mL each). The
aqueous layer was extracted with EtOAc (2 x 5 mL). The
combined organic extracts were dried (Na₂SO₄) and
concentrated in *vacuo* to give title compound as an orange
20 yellow-oil (81 mg crude).

B.




25

A solution of the Part A compound (70 mg; 0.17
mmol), tert-butyl bromoacetate (66 mg; 0.34 mmol), and
iPr₂NEt in DMF (0.5 mL) was stirred at RT for 2 days.
LC/MS showed that the reaction was complete and clean.

The crude reaction mixture was partitioned between H₂O (30 mL) and EtOAc (20 mL). The aqueous layer was extracted with Et₂O (2 x 10 mL); the combined organic extracts were dried (MgSO₄) and concentrated *in vacuo* to give the crude amino-*tert*-butyl ester.

This crude product was stirred in a 1:1 solution of CHCl₃ and TFA (2 mL) for 18 h at RT. The solution was then concentrated *in vacuo* and purified by preparative reverse-phase HPLC (as in Example 10). The purified material was lyophilized from MeOH-H₂O to give the title compound (71 mg; 71%) as a white lyophilate. [M + H]⁺ = 471.2

Example 126




15

The title compound was synthesized following the same procedure as described above in Example 125 except that (S)- α -methyl benzylamine was replaced by (R)- α -methyl benzylamine in the synthesis of the part A compound. The title compound was obtained in 67% yield (66 mg) overall. [M + H]⁺ = 471.2

25


Example 127



A mixture of Example 7 Part A compound (30 mg, 0.098 mmol), D-alanine *tert*-butyl ester hydrochloride (23 mg; 0.127 mmol), Et₃N (5 drops) and 4A molecular sieves

in MeOH (2 mL) was stirred at RT for 4 h. NaBH₄ (12 mg, 0.0294 mmol) was added and the reaction was stirred at RT for 30 min. The reaction mixture was then concentrated in vacuo, diluted with CH₂Cl₂ (2 mL), and filtered through cotton. TFA (1 mL) was added to the filtrate and the reaction was stirred at RT overnight. The reaction mixture was concentrated in vacuo, diluted with EtOAc, washed several times with sat'd. aqueous NaHCO₃, then with brine. The organic phase was dried (MgSO₄) and concentrated in vacuo. The residue was purified by preparative HPLC (YMC ODS 30mm x 250mm reverse-phase column; flow rate = 25 mL/min; 30 min continuous gradient from 50:50 A:B to 100% B, where A = 90:10:0.1 H₂O:MeOH:TFA and B = 90:10:0.1 MeOH:H₂O:TFA) to provide the title compound (7.8 mg, 21%) as a white lyophilate. [M + H]⁺ = 381.1


Example 128



20

Title compound (20% overall yield) was synthesized using the same procedure as described in Example 125, using D-phenylalanine tert-butyl ester hydrochloride instead of D-alanine tert-butyl ester hydrochloride. [M + H]⁺ = 457.2

Example 129



30

15


CC1=C(C2=CC=CC=C2O2)N=C(C3=CC=CC=C3)C1CCOC4=CC=CC=C4CN(Cc5ccc(Oc6ccccc6)cc5)C(C7=CC=CC=C7)C(=O)O

30

30


Examples 131 to 135

Other analogs in this series were prepared by analogous procedures and are shown in the following table:




5

Example No.	R ^{3c}	[M+H] ⁺
131	(S) -CH ₃	563.2
132	 (S)	639.3
133	 (R)	591.4
134	 (R)	579.3
135	 (R)	635.4


Example 136

A.



5

A solution of the secondary amine ethyl ester (72 mg; 0.183 mmol)




10

(prepared as described in Example 3 Part A) in MeOH (2 mL) and aqueous NaOH (2 mL of a 1M solution) was heated under reflux for 12 h. The pH of the solution was adjusted to 5 (with aqueous 1M NaOH and 1M HCl), upon which a colorless solid precipitated. This was filtered off and the filtrate was extracted with EtOAc (3x); the combined organic extracts were dried (Na₂SO₄) and concentrated *in vacuo* to give the crude title amino acid as a colorless solid (97 mg).

20

B.



To a solution of the Part A amino acid (15 mg; 0.04 mmol) in dioxane:H₂O (1:1, 8 mL) was added K₂CO₃ (22 mg;

5

Example 137



mg; 0.18 mmol)



20

25

30

5

COC1=CC=C(OC(=O)CN(C1)Cc2ccc(OCCc3c(C)c(Ph)n3)cc2)C=C1

A solution of 2-methoxyphenol (2 g, 16.1 mmol), N,N-dimethylaniline (1.95 g, 16.1 mmol), phosgene (8.34 mL of a 1.93 M solution in toluene, 16.1 mmol) and a catalytic amount of DMF in chlorobenzene (5 mL) was stirred in a pressure tube for 2 h at 80°C. The organic layer was separated and concentrated in vacuo. The residue was distilled (Buchi Kugelrohr; bp = 115°C @ 10 mm Hg) to provide 2-methoxyphenyl chloroformate (1.5g; 50%) as a clear oil.

COC1=CC=C(OC(=O)CN(C1)Cc2ccc(OCCc3c(C)oc(c3)Ph)cc2)C=C

25

CCOC(=O)NCc1ccc(OCCc2c(C)c(oc2=Nc3ccccc3)cc1)cc1

(prepared as described in Example 7 Part B),
2-methoxyphenyl chloroformate (8 mg, 0.05 mmol; prepared
as above) and polyvinylpyridine (Aldrich; 16 mg, 0.3
mmol) in CH_2Cl_2 (1 mL) was stirred for 30 min at RT.

5 Amine resin WA21J (Supelco; 200 mg) was added and the
mixture was stirred at RT for 30 min in order to remove
unreacted chloroformate. The reaction mixture was
filtered and concentrated in vacuo to give the desired 2-
methoxyphenyl carbamate-ester.

10 The ester was treated with a solution of 30% TFA in
 CH_2Cl_2 (5 mL) overnight. Volatiles were removed in vacuo
to give the crude acid. This material was purified via
solid-phase extraction using an anion exchange column
(CHQAX13M6 column; United Technologies; 3 g of sorbent in
15 a 6 mL column) by the exemplary procedure outlined below.

1) The column was conditioned with MeOH (10 mL) and CH_2Cl_2
(10 mL).

20 2) The crude acid was dissolved in a minimal volume of
 CH_2Cl_2 and loaded onto the SAX column.

3) The cartridge was washed with CH_2Cl_2 (10 mL),
 $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10 mL of a 4:1 $\text{CH}_2\text{Cl}_2:\text{MeOH}$ solution).

25 4) The product was eluted with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (10 mL of a 4:1
 $\text{CH}_2\text{Cl}_2:\text{MeOH}$ solution).

The product-containing fractions were concentrated
30 in vacuo on a Speed Vac to afford title compound as an
oil. Analytical reverse-phase HPLC (standard conditions)
indicated that the purity of the product was 90%. In
addition LC/MS gave the correct molecular ion $[(\text{M}+\text{H})^+ =$
517.3] for the desired title compound.


35

CC1=C(C(=O)N(C1)C2=CC=CC=C2OCC3=CC=CC=C3)C(=O)N(C2)C4=CC=CC=C4C(=O)OCCOC(=O)CN(Cc1ccc(OCC2=C(C)N(C2=O)C3=CC=CC=C3)cc1)C(=O)Cl

5

CCOC(=O)CNCCc1ccc(OCCc2c(C)c(Oc3ccccc3)n2)cc1

B.



20

- 128 -

A mixture of the Part A carbamoyl chloride (20 mg; 0.045 mmol), 3,5-dichlorophenol (16 mg; 0.07 mmol), and pyridine (0.5 ml) was stirred at 80°C for 16 h. Pyridine was removed *in vacuo* and the residue was purified via
5 solid-phase extraction using a CHQAX1 cartridge (2 g of sorbent in a 6 ml column, 0.3 mg/g) by the procedure outlined below:


- 1) The column was conditioned with MeOH (10 mL) and
10 CH₂Cl₂ (20 mL)
- 2) The reaction mixture in CH₂Cl₂ was loaded onto the SAX column
- 15 3) The product was eluted with CH₂Cl₂ (10 mL)

The product-containing fractions were concentrated *in vacuo* using a Speed Vac over 16 h to afford the pure aryl carbamate-tert-butyl ester which was treated with a
20 solution of 30% TFA in CH₂Cl₂ overnight. Volatiles were removed using a Speed Vac for 16 h to afford the crude acid final product. The product was initially purified via solid-phase extraction using a Varian SAX cartridge (2 g of sorbent in a 6 mL column, 0.3 meq/g) by the
25 procedure outlined below:


- 1) The column was conditioned with MeOH (10 mL) and CH₂Cl₂ (20 mL)
- 30 2) The reaction mixture in CH₂Cl₂ was loaded onto the SAX column
- 3) The column was rinsed with CH₂Cl₂ (10 mL)
- 35 4) The column was rinsed with 10% MeOH in CH₂Cl₂ (10 mL)

The product-containing fractions were concentrated *in vacuo* using a Speed Vac for 16 h to afford the purified product (20 mg, 80%) as a solid. Reverse phase HPLC analysis (YMC S5 ODS 4.6 x 33 mm column, continuous gradient from 50% A to 100% B for 2 min at a flow rate of 5 mL/min [Solvent A = 10%MeOH/90%H₂O/0.2% H₃PO₄; Solvent B = 90%MeOH/10% H₂O/0.2% H₃PO₄]) indicated that the product purity was 96%. In addition, LC/MS gave the correct molecular ion [(M+H)⁺ = 555.2] (electrospray) for the title compound.

15



A.




- 130 -

5

COC1=CC=C(C=C1)COCC(=O)N(C)Cc2ccc(OCCc3c(C)c(oc3=Nc4ccccc4)cc5)cc2

The title compound was prepared as part of a solution phase library which was run using the following standard procedure.

15



20

25


- 1) The column was conditioned with MeOH (10 mL) and CH₂Cl₂ (10 mL)

3) The cartridge was washed successively with CH_2Cl_2 (10
5 mL), 20% $\text{MeOH}/\text{CH}_2\text{Cl}_2$ (10 mL).


4) The product was eluted with a solution of 20% MeOH/CH₂Cl₂ (10 mL).

10 The product-containing fractions were concentrated
in vacuo using a Speed Vac to afford the title compound.
Reverse Phase HPLC analysis using standard conditions
indicated that the product purity was 90%. In addition,
LC/MS (electrospray) gave the correct molecular ion
15 $[(M+H)^+ = 531.3]$ for the desired title compound.

20




A.



30


separated and concentrated *in vacuo* to give the crude title aryl chloroformate as crystals (2 g crude product).

B.




5

To a mixture of the Part A chloroformate (184 mg, 0.70 mmol) in CH₂Cl₂ (5 mL) and polyvinylpyridine (Aldrich; 315 mg, 1 mmol) was added a solution of the amino-tert-butyl ester (280 mg, 0.66 mmol)



(prepared as described in Example 7 Part B),
in CH₂Cl₂ (5 mL). The reaction was stirred at RT for 15
15 min. Resin-bound amine (WA21J, Supelco; 150 mg) was
added to the mixture. The reaction mixture was stirred
for another 15 min. The resin-bound amine and
polyvinylpyridine were filtered off and the filtrate was
concentrated *in vacuo* to give the crude product. The
20 crude product was chromatographed (SiO₂; hexane/EtOAc
1:4) to provide title compound (0.30 g, 70%).

C.



25

A solution of Part B compound (75 mg; 0.42 mmol) in 20 ml MeOH was hydrogenated in the presence of 20 mg of 10% Pd/C under an atmosphere of H₂ (balloon) for 24 h.

5

CC1=C(C2=CC=CC=C2O1)N=C(C3=CC=CC=C3)C1=CC=CC=C1CCOC1=CC=C(C=C1)CN(C(=O)O)C(=O)OC1=CC=C(C=C1)OC(F)F


30

Following the Example 141 procedure, the analogous compound was prepared [(M+H)⁺ = 553.2]:



5

Example 144




Example 143: $[M + H]^+ = 593.4$


10














Examples 145 to 305















15













Table 4














Example No.	R ^{3d}	[M+H] ⁺
145		487.2

Example No.	R ^{3d}	[M+H] ⁺
146		539.3
147		593.2
148		449.3
149		501.3
150		517.2
151		532.2
152		589.3
153		546.3
154		532.2
155		579.2
156		593.2
157		593.3
158		579.2

Example No.	R ^{3d}	[M+H] ⁺
159		579.2
160		531.2
161		527.2
162		525.2
163		515.2
164		529.2
165		527.2
166		519.3
167		517.3
168		547.3
169		577.3
170		531.3
171		531.3
172		545.3


Example No.	R ^{3d}	[M+H] ⁺
173		531.3
174		571.2
175		567.3
176		547.3
177		545.3
178		579.2
179		591.2
180		535.2
181		505.2
182		521.1
183		566 + 588
184		571.1

- 139 -

Example No.	R ^{3d}	[M+H] ⁺
195		529.4
196		531.3
197		561.3
198		555.3
199		513.3
200		557.4
201		543.3
202		571.4
203		572.3
204		541.3
205		543.4















Example No.	R ^{3d}	[M+H] ⁺
218		573.1
219		519.2
220		535.2
221		585.2
222		519.2
223		535.2
224		585.2
225		561.2














Table 5
























5












Example No.	R ^{3d}	[M+H] ⁺
226		545.2













Example No.	R ^{3d}	[M+H] ⁺
227		593.1
228		449.2
229		501.2
230		517.2
231		532.2
232		487.3
233		546.3
234		532.2
235		579.2
236		593.2
237		593.3
238		579.2
239		579.2
240		531.2

Example No.	R ^{3d}	[M+H] ⁺
241		527.2
242		525.2
243		515.2
244		529.2
245		527.2
246		517.3
247		517.3
248		547.3
249		577.3
250		543.1
251		531.3
252		545.3
253		531.3

Example No.	R ^{3d}	[M+H] ⁺
254		571.2
255		567.3
256		547.3
257		545.3
258		593.4
259		503.2
260		579.2
261		505.2
262		521.1
263		566/567

Example No.	R ^{3d}	[M+H] ⁺
264		571.1
265		505.2
266		521.1
267		566/567.0
268		523.3
269		501.3
270		539.2
271		529.3
272		549.2
273		523.2
274		513.3

Example No.	R ^{3d}	[M+H] ⁺
275		519.2
276		539.2
277		547.3
278		552.3
279		541.3
280		563.3
281		555.3
282		543.3
283		529.3
284		515.3
285		515.3


Example No.	R ^{3d}	[M+H] ⁺
286		515.3
287		535.2
288		529.2
289		577.3
290		515.2
291		529.2
292		527.3
293		537.3
294		531.3
295		555.2
296		571.3
297		573.2

Example 149



5

To a solution of the secondary amine-ester (2.1 g;
5.52 mmol)




[M + H]⁺ = 501.2; ¹H NMR (400 MHz; CDCl₃): δ 7.93-7.99 (m, 2H), 7.38-7.43 (m, 3H), 7.23 (q, 1H, J = 8 Hz), 7.02-7.12 (m, 3H), 6.98-7.02 (m, 2H), 6.82-6.89 (m, 2H), 4.71 (s, 1H), 4.64 (s, 1H), 4.25 (t, 2H, J = 7 Hz), 4.07 (s, 2H), 2.90-2.98 (m, 2H), 2.37 (s, 3H), 2.29 (s, 3H)

Example 230



To a 0°C solution of the secondary amine (3.0 g; 7.9 mmol)



in CH₂Cl₂ (45 mL) were successively added pyridine (0.8 mL; 9.9 mmol) and 4-methoxyphenyl chloroformate (1.3 mL; 8.7 mmol). The reaction was stirred at 0°C for 3 h, at which point starting material had been consumed (by analytical HPLC). The reaction solution was washed with aqueous HCl (2 x 25 mL of a 1 M solution), brine (2x), dried (Na₂SO₄), and concentrated *in vacuo*. The crude product was chromatographed (SiO₂; stepwise gradient from 4:1 to 3:7 hex:EtOAc) to provide the desired carbamate-ester (4.2 g; 100%). The ester was dissolved in THF:MeOH:H₂O (50 mL of a 3:1:1 solution) and LiOH.H₂O (0.5 g; 11.9 mmol) was added. The solution was stirred overnight at RT. Starting material was still present by HPLC. More LiOH.H₂O (0.2 g; 4.8 mmol) was added and the mixture was briefly heated to solubilize the LiOH, then stirred at RT for 4 h. The reaction was complete at this point, and the mixture was acidified to pH 3 with excess aqueous 1 M HCl, then organic solvents were removed in *vacuo*. The residual aqueous phase was extracted with EtOAc (3 x 50 mL). The combined organic extracts were successively washed with H₂O and brine (50 mL each), dried (Na₂SO₄), filtered and concentrated *in vacuo* to give title compound as a colorless solid (3.07 g; 75%).

[M + H]⁺ = 517.2; ¹H NMR (400 MHz; CDCl₃): δ 7.96–7.98 (m, 2H), 7.4–7.45 (m, 3H), 7.2–7.3 (m, 2H), 7.0–7.1 (m, 2H), 6.8–7.0 (m, 4 H), 4.65 (s, 1H), 4.55 (s, 1H), 4.20–4.24 (m, 2H), 4.02 (s, 2H), 3.77 (s, 3H), 3.00 (s, 2H), 2.38 (s, 3H).

5

COc1ccc(cc1)OC(=O)NCCN(Cc2ccc(OCCc3c(C)c(C4=CC=CC=C4)on3)cc2)C(=O)O

15

CC1=C(C(=O)O)C(=O)N(C1)Cc2ccc(OCC3=C(C)C(=O)N(C3)C4=CC=CC=C4)cc2

25

COC(=O)c1ccc(OCC2=C(C)N(c3ccccc3)C2=O)cc1

5

10



15

20




25

CC1=C(C2=CC=CC=C2)OC(=N1)C3=CC=CC=C3CCOC4=CC=C(C=C4)CN(C(=O)OC5=CC=C(C=C5)Br)CC(=O)O

10 Hz, 2H)


CC1=C(C2=CC=CC=C2O2)N(C3=CC=CC=C3)C1CCOC4=CC=CC=C4CN(C(=O)O)C(=O)Oc5ccc(OCF2)c(OC)c5

A.




30

B.




- 5 To a mixture of Part A chloroformate (50 mg, 0.237 mmol) and polyvinylpyridine (PVP) (75 mg, 0.70 mmol) was added a CH_2Cl_2 solution (2 mL) of the amino-tert-butyl ester (100 mg, 0.237 mmol),



- 10 (prepared as described in Example 7 Part B).


The reaction was stirred at RT for 15 min. Resin-bound amine (WA21J, Supelco; 150 mg) was added to the mixture. The reaction mixture was stirred for another 15 min. The Resin-bound amine and PVP were removed via
 15 filtration and the filtrate was concentrated *in vacuo* to give the crude product. The crude product was chromatographed (SiO_2 ; hexane/EtOAc 1:4) to provide title compound (0.1 g, 70%).

20 C.



- 25 A solution of the Part B phenol-tert butyl ester compound (60 mg; 0.10 mmol), Bu_4NBr (0.32 mg, 0.001 mmol), aqueous NaOH (0.5 mL of a 1 M solution; 0.5 mmol) and isopropanol (1 mL) in a pressure tube was cooled to -50°C . Freon gas was bubbled into the solution for 1 min. The tube was sealed and heated to 80°C for 12 h.



The mixture was extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated *in vacuo* to give the crude difluoromethoxy ether-tert butyl ester as an oil. The crude ester was then treated with a solution of 30% TFA in CH_2Cl_2 overnight. Volatiles were removed in *vacuo* and the residue was purified using preparative reverse-phase HPLC (as in Example 127, except that the continuous gradient used was from A:B 70:30 to 100% B) to afford two products, the desired title difluoromethoxy ether-acid (13 mg; 23%) and the phenol-acid set out below (32 mg; 63%). Reverse phase HPLC analysis using standard conditions indicated that the product purity was >92%. In addition LC/MS (electrospray) gave the correct molecular ion $[(M+H)^+ = 553.2 \text{ and } 503.2 \text{ respectively}]$ for the two compounds.

**Phenol-Acid**

20

Examples 307 and 308

Following the above general procedure of Example 306, the following compounds were prepared:

**Example 307****Example 308**25 Example 307: $[M + H]^+ = 553.2$ Example 308: $[M + H]^+ = 503.2$

CC1=C(C(=N1C2=CC=CC=C2)OCCOC3=CC=C(C=C3)CN(CS(=O)C4=CC=CC=C4)CC(=O)O)C


To a mixture of phenyl chlorothionoformate (11 mg, 0.063 mmol) and triethylamine (6.5 mg, 0.063 mmol) was added a solution of the amino-tert-butyl ester (20 mg, 0.053 mmol),




15


25

25


$$[M + H]^+ = 503.2$$

Example 311

5 To a mixture of the amine-tert butyl ester (306 mg, 0.73 mmol)



(prepared as described in Example 7 Part B),
and p-phenoxybenzoic acid (220 mg; 1.02 mmol; 1.4 equiv)
10 in CH₃CN (20 mL) was added BOP reagent (372 mg, 0.84 mmol, 1.15 equiv) in a single portion followed by iPr₂NEt (0.5 mL; 2.9 mmol; 3.9 equiv) dropwise. The reaction was stirred overnight at RT, after which volatiles were removed *in vacuo*. The residue was dissolved in EtOAc and
15 washed with aqueous 1N HCl. The aqueous layer was extracted with EtOAc (2x) and the combined organic extracts were washed with H₂O, sat'd aqueous NaHCO₃ and brine, dried (Na₂SO₄) and concentrated *in vacuo* to give the desired product. The resulting crude amide-ester was
20 used in the next step without further purification.

A solution of the crude amide ester in 40% TFA-CH₂Cl₂ (25 mL) was stirred for 5h at RT. Volatiles were removed *in vacuo* and the crude acid was purified by Prep HPLC (YMC S5 ODS 30mm x 250 mm reverse phase column; flow rate
25 = 25 mL/min; 30 min continuous gradient from 70:30 A:B to 100% B; solvent A = 90:10:0.1 H₂O:MeOH:TFA; solvent B = 90:10:0.1 MeOH:H₂O:TFA) to yield title compound (238 mg; 58% yield over 2 steps) as a white solid. Analytical Reverse-phase HPLC: Retention time = 7.53 min.
30 (Continuous gradient solvent system: from 50% A:50% B to


5

CC1=C(COP(=O)(OC(=O)N)OC(=O)c2ccc(Oc3ccccc3)cc2)OC(=O)c4ccccc4N1

10


CCOC(=O)CNCCc1ccc(OCCc2c(C)c(Oc3ccccc3)n2)cc1

30

Example 312

5

- 1) To a solution of the secondary amine-tert-butyl ester (25 mg; 0.06 mmol)



- (prepared as described in Example 7 Part B),
10 in THF (0.5 mL) was added 2-naphthalene carboxylic acid (25 mg; 0.15 mmol; 2.5 equiv).

2) HOAT (48 mg; 0.35 mmol; 5.8 equiv) was added.

15 3) DMF (50 μ L) was added.

4) EDCI ((20 mg, 0.1 mmol, 1.8 m eq) was added.

5) The reaction vessel was shaken for 24 h at RT.


20

6) The reaction was diluted with MeOH (2 mL) and filtered.


7) The amide-tert butyl ester was purified by preparative
25 HPLC (YMC S5 ODS 20 x 100 mm column; flow rate = 25 mL/min; 10 min continuous gradient from 70:30 A:B to 100% B; solvent A = 90:10:0.1 H₂O:MeOH:TFA; solvent B = 90:10:0.1 MeOH:H₂O:TFA).

8) The fractions containing the purified amide-ester were treated with a solution of TFA in CH_2Cl_2 (0.5 mL of a 1:1 solution) overnight. The reaction was concentrated in vacuo (Speed Vac) to give title compound (8 mg; 25%). Reverse-phase analytical HPLC showed that the purity of the product was > 88%; LC/MS (electrospray detection) gave the correct $[\text{M} + \text{H}]^+ = 521.2$ for the title compound.

Example 313



A mixture of the amino-ester (20 mg; 0.0474 mmol),



(prepared as described in Example 7 Part B),


thiophene-2-carboxylic acid (9.1 mg, 0.71 mmol), EDCI (26 mg, 1.4 mmol) and DMAP (a catalytic amount) was dissolved in CH₂Cl₂ (2 mL) and stirred at RT overnight. The

reaction solution was successively washed with aqueous 1N HCl (2 mL) and sat'd aqueous NaHCO₃ (2 mL). To the organic phase was then added 0.5 g anhydrous Na₂SO₄, and 0.2 g WA21J amine-bound resin (Supelco). The mixture was shaken for 0.5 h and the solids were filtered off. TFA (2.0 mL) was added to the filtrate and the solution was shaken at RT overnight. The reaction solution was concentrated in vacuo using a Speed Vac for 16 h to afford title compound as a yellow oil. Reverse phase analytical HPLC (YMC S5 ODS 4.6 x 33 mm column, continuous gradient from 100% A to 100%B for 2 min at a

5


Example 314

10



15

mmol),



20


25

10

Examples 315 to 391

Following one of the above procedures, the following compounds in Tables 6 and 7 of the invention were prepared.

Table 6: (Amide-Acids)




Example No.	R ^{3e}	[M+H] ⁺
315		521.2
316		507.3
317		563.1
318		561.2
319		499.3
320		559.2
321		491.1
322		522.2
323		491.2
324		543.3












- 166 -

- 167. -


Table 7: (Amide-Acids)




Example No.	R ³	[M+H] ⁺
362		499.2
363		547.2
364		563.2
365		561.1
366		595.1
367		593.1
368		595.1
369		597.1

Example No.	R ³	[M+H] ⁺
370		563.1
371		547.2
372		563.1
373		577.2
374		561.2
375		561.2
376		536.2
377		577.2
378		577.2
379		615.3
380		499.3

- 170 -


Example 392

A.



5


To a solution of the amine (47 mg; 0.12 mmol)



10

(prepared as described in Example 3 Part A),
 in CH_2Cl_2 (5 mL) were added $i\text{Pr}_2\text{NEt}$ (0.1 mL; 0.57 mmol)
 and DMAP (14 mg; 0.12 mmol) followed by benzyl isocyanate
 (24 mg; 0.18 mmol). The reaction was stirred for 14h,
 15 then passed through an SCX cartridge [the 3g SCX
 cartridge was prewashed successively with MeOH (10 mL)
 and CH_2Cl_2 (5 mL)] by eluting with CH_2Cl_2 (15 mL). The
 filtrate was concentrated in vacuo to give the crude urea
 Part A compound (53 mg; 84%), which was sufficiently pure
 20 to be used in the next step without further purification.

B.



25


10

Cc1c(CCOc2ccc(CN(Cc3ccccc3)c4cc(OC)cc4)cc2)c5ccccc5n1COC(=O)CN(Cc1ccc(OCC2=C(C)OC(=N2)c3ccccc13)cc1)C(=O)Nc2ccc(OC)cc2COC(=O)CNc1ccc(OCCc2c(C)c(Oc3ccccc3)n2)cc1

20

25

B.




A solution of Part A compound (0.14 g, 0.26 mmol) and LiOH (0.1 g, 4.3 mmol) in H₂O/THF (5 ml of a 40:60 solution) was stirred for 12 h at 25°C. The reaction mixture was acidified with HOAc and extracted with EtOAc (2x). The combined organic extracts were dried (MgSO₄) and concentrated in vacuo to provide title compound (0.12 g; 90%) as a colorless oil. [M + H]⁺ = 516

15


¹H NMR (CD₃OD; δ): 7.94 (m, 2H), 7.45 (m, 3H), 7.23 (m, 3H), 6.80 (m, 2H), 6.80 (m, 3H), 4.58 (s, 2H), 4.23 (t, J = 7.9 Hz, 2H), 3.81 (s, 2H), 3.73 (s, 3H), 2.98 (t, J = 7.9 Hz, 2H), 2.36 (s, 3H).

20


Example 394



A.




5 A solution of the previously described carbamoyl chloride (Example 139 Part A compound; 0.15 g; 0.34 mmol)



10 N-methyl-p-anisidine (0.14 g, 1.0 mmol) and K_2CO_3 (0.15 g, 1.1 mmol) in 5 ml of acetone was stirred at 25 °C for 12 h. The reaction mixture was concentrated in vacuo to yield an oily residue, which was chromatographed (SiO_2 ; 1.5% MeOH/ CH_2Cl_2) to provide title compound (0.12 g; 65%) as a colorless oil.

15 B.



20 A solution of Part A compound (0.12 g, 0.22 mmol) and LiOH (0.050 g, 2.1 mmol) in H_2O /THF (5 mL of a 40:60 solution) was stirred at RT for 12 h. The reaction mixture was concentrated in vacuo to yield an oily residue, which was purified by preparative HPLC (YMC S5 ODS 30 x 250 mm column; flow rate = 25 ml/min. 30 min
25 continuous gradient from A:B = 50:50 to 100%B; solvent

A=90:10:0.1 H₂O:MeOH:TFA; solvent B = 90:10:0.1

MeOH:H₂O:TFA) to provide title compound (59 mg, 50%) as a colorless oil. $[M + H]^+ = 530.3$

5 NMR (CDCl₃): 7.99 (d, 6.2 Hz, 2H, 7.45 (m, 3H), 7.24 (m, 3H), 6.82 (d, 6.2 Hz, 2H), 6.79 (m, 1H), 6.63 (m, 1H), 6.55 (s, 1H), 4.24 (s, 2H), 4.16 (t, 7.8 Hz), 2H), 3.72 (s, 3H), 3.59 (s, 2H), 3.16 (s, 2H), 3.02 (t, 7.8 Hz, 2H), 2.40 (s, 3H).


10




Examples 395 to 410

Utilizing one of the above procedures, the analogs in Tables 8 and 9 were synthesized.

15

Table 8: (Urea-Acids)



Example No.	R ^{3f}	[M+H] ⁺
395		562.3
396		546.3
397		554.2














Example No.	R^{3f}	$[M+H]^+$
398		532.3
399		522.3
400		546.3
401		516.3


Table 9: (Urea-Acids)



5

Example No.	R^{3f}	$[M+H]^+$
402		562.3
403		546.3
404		554.2

Example No.	R ^{3f}	[M+H] ⁺
405		532.3
406		522.3
407		546.3
408		516.3
409		516.3


Example 410

5


The title compound was prepared as part of a
solution phase library run using the following procedure:

10

A.



5 To a mixture of 1-naphthylsulfonyl chloride (26.8 mg, 0.12 mmol) and DMAP (2 mg, 0.016 mmol) in pyridine (2 mL) was added a solution of the amino-t-butyl ester



(prepared as described in Example 8)

10 (20 mg, 0.05 mmol) in pyridine (0.6 mL). The reaction was stirred at RT for 20 h. Resin-bound amine (WA21J, Supelco; 5.8 mmol/g loading; 150 mg) was added to the mixture. The reaction was stirred for a further 4 h. The resin was filtered off and the filtrate was


15 concentrated *in vacuo* to give the crude product, which was chromatographed (CUSIL12M6 column; United technology; 2 g of sorbent in a 6 mL column) by the procedure outlined below.

20 1) The column was conditioned with hexane (20 mL).

2) The residue was dissolved in a minimal volume of EtOAc and loaded onto the silica gel column.


25 3) The cartridge was eluted with Hex/EtOAc(3:1),
Hex/EtOAc (1:1). The desired fraction (identified by
TLC) was collected and concentrated to give title
compound as a viscous oil which was used in the next step
without any further purification.

B.




- 5 Et₃N ((0.3 ml of a 1M solution in CH₂Cl₂) and TMSI
(0.3 ml of a 1M solution in CH₂Cl₂) were successively
added to a solution of Part A compound in CH₂Cl₂. The
reaction mixture was stirred at RT for 12h and then was
concentrated *in vacuo* to give the crude product. The
10 product was purified by solid-phase extraction using a
CHQAX12M6 column (United technology; 2 g of sorbent in a
6 mL column) by the procedure outlined below.
- 1) The column was conditioned with CH₂Cl₂ (25 mL).
- 15 2) The residue was dissolved in a minimal volume of CH₂Cl₂
and loaded onto the SAX column.
- 3) The cartridge was washed successively with CH₂Cl₂ (25
20 mL), CH₂Cl₂/MeOH (5% MeOH, 15 mL), CH₂Cl₂/MeOH (50% MeOH,
15 mL), MeOH (20 mL).
- 4) The product was eluted with a solution of 1% TFA in
MeOH (20 mL).
- 25 The final product-containing fraction was collected
and concentrated *in vacuo* using a Speed Vac to afford
BMS-329075 (16 mg; 62%). Reverse-phase analytical HPLC
indicated that the product purity was 90%. In addition,
30 LC/MS (electrospray) gave the correct molecular ion
[(M+H)⁺ = 557.1] for the desired compound.

Example 411



A.




(X = halogen, alkyl, CF₃, CF₃O, etc.)


10 The following general procedure was utilized for the
preparation of the requisite substituted benzyl sulfonyl
chlorides:

Cl₂ gas was bubbled into a 0°C solution of 4-fluorobenzyl mercaptan (1.0 g, Lancaster) in glacial acetic acid (100 mL) and H₂O (5.0 mL) for 1h. The reaction mixture was then poured into ice-H₂O and immediately extracted with CH₂Cl₂ (200 mL); the organic phase was cautiously washed successively with H₂O (200 mL), aqueous saturated NaHCO₃ (2 x 100 mL), and finally brine (200 mL). The organic phase was dried (MgSO₄) and concentrated in vacuo to furnish 4-fluorobenzyl sulfonyl chloride as a colorless solid (1.3 g; 89%).

25 B.



To a solution of the secondary amine methyl ester
(25 mg; 0.066 mmol)



5 (prepared as described in Example 6),
in pyridine (0.8 mL) was added 4-fluorobenzyl sulfonyl
chloride (68 mg; 0.33 mmol; 5 equiv). The mixture was
heated to 75°C, stirred overnight at 75°C, and then
concentrated in vacuo. The black residue was treated
10 with aqueous LiOH (1.0 mL of a 0.3 M solution) in
H₂O/MeOH/THF for 18 h, then concentrated in vacuo. The
residue was acidified with 1.0 M aqueous HCl to pH = 1-2
and extracted with EtOAc (2x), dried (Na₂SO₄) and
concentrated in vacuo to give the crude product.
15 Purification by preparative HPLC (YMC S5 ODS 20mm x 250
mm reverse-phase column; 15 min continuous gradient from
60:40 A:B to 100% B with 10 min hold time, where A =
90:10:0.1 H₂O:MeOH:TFA and B = 90:10:0.1 MeOH:H₂O:TFA;
flow rate = 25 mL/min) gave the title compound (12 mg;
20 34%) as a white solid. [M + H]⁺ (LC/MS) = 539.1

Examples 412 to 456

Utilizing one of the above procedures, the analogs in Tables 10 and 11 were synthesized.













Example No.	R ^{3g}	[M+H] ⁺
423		535.3
424		539.1
425		541.2
426		589.0
427		573.2
428		555.2
429		555.3
430		589.2
431		535.3
432		605.3
433		577.4

Table 11: (Sulfonamide-Acids)




Example No.	R ^{3g}	[M+H] ⁺
434		549.4
435		557.3
436		506.3
437		549.3
438		541.2
439		521.3
440		533.3
441		535.4
442		575.3
443		678.3

- 185 -

CC1=C(C(=N1C(=O)C2=CC=CC=C2)COC3=CC=NC=C3CN(CCC(=O)O)Cc4ccc(Oc5ccccc5)cc4)CCOC(=O)c1ccc(OCCc2c(C)c(oc2=Nc3ccccc3)cn1)cc1

5

B.




20

- 186 -


vacuo to furnish title compound as an oil. This crude material was used in the next reaction without further purification.

5 C.



To a -78°C solution of oxalyl chloride (0.22 mL, 2.6 mmol) and DMSO (0.37 mL, 5.2 mmol) in CH_2Cl_2 (15 mL) was added dropwise a solution of Part B compound (0.42 g of crude material in 5 mL CH_2Cl_2). The reaction mixture was stirred for 2 h at -78°C and then Et_3N (1 mL) was added dropwise. The reaction mixture was stirred for an additional 0.5 h at -78°C and then was slowly warmed to 25°C . The reaction mixture was diluted with EtOAc (200 mL) and washed successively with aqueous NaHCO_3 and brine. The organic layer was dried (MgSO_4), then concentrated in vacuo to provide title compound (0.40 g; 95%) as an oil, which was used in the next step without further purification.

D.




25

A mixture of Part C compound (<0.82 mmol), glycine methyl ester hydrochloride (0.5g., 4.0 mmol), $\text{NaBH}(\text{OAc})_3$ (0.85g., 4.0 mmol) and DCE (10 mL) was stirred at 25°C for 12 h. The reaction mixture was then diluted with EtOAc (50 mL) and washed successively with aqueous NaHCO_3 and brine. The organic layer was dried (MgSO_4), then concentrated in vacuo to give title compound (0.31 g;

82%) as an oil (>95% pure by analytical reverse-phase HPLC) which was used in the next step without further purification.

5


E.



A mixture of Part D compound (0.050 g; 0.13 mmol),
10 4-phenoxybenzaldehyde (0.048 g; 0.26 mmol), NaBH(OAc)₃
(0.082 g; 0.39 mmol) in DCE (10 mL) was stirred at 25°C
for 12 h. The reaction mixture was diluted with EtOAc
(50 mL) and washed successively with aqueous NaHCO₃ and
brine. The organic layer was dried (MgSO₄), then
15 concentrated in vacuo to give the tertiary amino methyl
ester as an oily residue. To this residue, LiOH (0.050
g) and H₂O/THF (2 mL of a 60/40 solution) were added and
the reaction was stirred at RT for 12 h. Preparative
HPLC (YMC S5 ODS 30 x 250mm column - continuous gradient
20 over 30 min; flow rate = 25 mL/min from 30:70 A:B to
100% B; A = 90:10:0.1 H₂O:MeOH:CF₃CO₂H; B = 90:10:0.1
MeOH:H₂O:CF₃CO₂H) provided title compound (0.021 g; 30%)
as a TFA salt.


25 ¹H NMR (CDCl₃) δ: 8.18 (s, 1H), 7.94 (d, 6.6 Hz, 2H),
7.86 (d, 8.8 Hz, 1H), 7.45 (m, 3H), 7.34 (m, 3H), 7.14
(t, 7.4 Hz, 1H), 7.02-6.92 (m, 5H), 6.81 (t, 8.8 Hz, 1H),
4.51 (m, 6H), 3.59 (s, 2H), 3.06 (t, 6.2 Hz, 2H)

30


Example 458

COC(=O)c1cc(O)ccn1

15 B.




C.




5

15




20

E.




5 To a solution of Part D compound (0.90g; 2.7 mmol),
glycine methyl ester hydrochloride (1.7 g, 13.5 mmol) in
1,2 dichloroethane (10 mL) was added NaBH(OAc)₃ (1.7 g,
8.1 mmol) in one portion. The resulting solution was
stirred at 25°C for 12 h. The reaction mixture was
10 concentrated in vacuo to give an oil, which was
chromatographed (SiO₂; 30% acetone in hexane) to provide
title compound (0.86 g; 83%) as a colorless oil.

F.



A solution of Part E compound (0.040 g, 0.1 mmol), 4-phenoxybenzaldehyde (0.030 g, 0.15 mmol) and $\text{NaBH}(\text{OAc})_3$ (0.060 g, 0.3 mmol) in DCE (10 mL) was stirred at RT for 12 h. The reaction mixture was concentrated in vacuo and the oily residue was chromatographed (SiO_2 ; 30% acetone in hexane) to provide the amino-ester title compound (56 mg; >95%) as a colorless oil.

G.



10

15

20




30


The reaction mixture was concentrated *in vacuo*, and the residue was chromatographed (SiO₂; stepwise gradient from 95:5 to 4:1 hex:EtOAc) to obtain title compound (1.43 g, 65%).

5

B.




- 10 A solution of Part A compound (1.01 g, 2.37 mmol) and TFA (8 mL) in CH₂Cl₂ (30 mL) was stirred at RT for 4.5 h. The solution was concentrated *in vacuo*, and the residue was dissolved in CH₂Cl₂ and filtered through a pad of solid K₂CO₃. The filtrate was concentrated *in vacuo* to give the corresponding crude amine. To a solution of the crude amine in THF (11.9 mL) were added pyridine (0.383 mL, 4.74 mmol) and 2,4-dinitrobenzenesulfonyl chloride (0.85 g, 3.19 mmol) and the solution was stirred at RT for 15h. Since some starting material still remained at this point, more sulfonyl chloride (0.32 g, 1.2 mmol) was then added. After a further 4 h, HPLC analysis indicated that all starting material had been consumed. The reaction mixture was diluted with Et₂O, washed with 1N aq HCl, saturated aq NaHCO₃ and brine, dried (MgSO₄), filtered and concentrated *in vacuo* to provide the crude 2,4-dinitrobenzenesulfonamide




To a solution of the crude 2,4-dinitrobenzene-sulfonamide in CH₃CN (3 mL) were added K₂CO₃ (excess) and tert-butyl bromoacetate (7.11 mmol). The reaction was stirred at RT overnight. HPLC analysis indicated the ratio of product to starting material was 2/1. More DMF (3 mL), K₂CO₃ and tert-butyl bromoacetate were added to the reaction mixture. The reaction was complete in 2 h. The reaction mixture was diluted with Et₂O, washed with 1N aq HCl, saturated NaHCO₃ and brine, dried (MgSO₄), and concentrated *in vacuo* to provide the crude tert-butyl ester. This crude material was chromatographed (SiO₂; hexanes/EtOAc; stepwise gradient from 9:1 to 2:1) to give title compound (0.663 g, 42% overall).

C.



To a solution of Part B compound (0.663 g, 0.995 mmol) in THF (2.5 mL) were added Et₃N (0.208 mL, 1.49 mmol) and mercaptoacetic acid (0.090 mL, 1.29 mmol). The reaction was stirred at RT overnight. The reaction mixture was then diluted with Et₂O, washed with 1N aq HCl, saturated NaHCO₃ and brine, dried (MgSO₄), and concentrated *in vacuo*. The residue was chromatographed (SiO₂; hexanes/EtOAc; stepwise gradient from 9:1 to 2:1) to give title compound (0.265 g, 61%).


D.



To a solution of Part C compound (0.015 g, 0.0344 mmol) in DCE (1 mL) were added 4-phenoxybenzaldehyde (0.103 mmol) and $\text{NaBH}(\text{OAc})_3$ (0.0365 g, 0.172 mmol). The reaction was stirred at RT for 15 h. The reaction mixture was filtered through a cotton plug to provide a clear solution, which was diluted with CH_2Cl_2 , washed with saturated aq NaHCO_3 and brine, dried (MgSO_4), and concentrated *in vacuo*. The crude product was purified by preparative HPLC (YMC S5 ODS 30x250 mm column: flow rate 25 mL/min, gradient 20%B to 100%B over 25 min, 100%B hold for 15 min, Retention time = 29.1 min) to furnish the tert-butyl ester. A solution of this material was dissolved in CH_2Cl_2 (1.3 mL) and TFA (0.5 mL) was added slowly. The reaction was stirred at RT overnight and was then concentrated *in vacuo*. The residue was then dissolved in CH_2Cl_2 , washed with H_2O , saturated aq NaHCO_3 and brine, dried (MgSO_4) and concentrated *in vacuo* to give title compound (0.012 g, 61%). LC/MS gave the correct $[\text{M} + \text{H}]^+ = 563.3$

Further analogs (as shown in the table below) were synthesized by the same reductive amination procedure as described in Example 459 Part D using Example 459 Part C compound and different aromatic aldehydes. In addition carbamate-acids such as Example 461 compound were also synthesized using the general method described previously for the synthesis of the Example 136 compound.


Table 12




Example No.	R ³	[M+H] ⁺
460		571.3
461		515.3
462		471.3

5

Example 463



A.



10

To a solution of the Example 230 acid (240 mg, 0.47 mmol) in DMF (2.0 mL) were added HOAT (68 mg, 0.49 mmol),
 15 EDAC (94 mg, 0.49 mmol) and 2-cyanoethylamine (34 mg, 0.49 mmol). The solution was stirred at RT for 18 h;


5
10

[illegible]

20
25
30

chromatographed (SiO₂; ether:CH₂Cl₂ 3:7) to furnish the title compound (100 mg; 53%) as a white solid.

C.




5


To a solution of Part B compound (100 mg, 0.17 mmol) in THF/1,4-dioxane (6:1, 1.4 mL) was added aqueous NaOH solution (0.6 mL of a 1.0 M solution, 3.5 equiv). The mixture was stirred at RT for 14 h and then acidified to ~pH 2 with 1.0 M aqueous H₃PO₄ solution. EtOAc (30 mL) was added, and the organic phase was washed with water (15 mL) and brine (15 mL), dried (MgSO₄) and concentrated in vacuo. The residue was chromatographed (SiO₂; 4% MeOH/CH₂Cl₂) to give the title tetrazole (35 mg; 38%) as a white foam. LC/MS (electrospray) gave the correct molecular ion: [M + H]⁺ = 541.3

20

Example 464




A.




25

15 B.




- 199 -

C.



- 5 To a solution of Part B compound (40 mg, 0.116 mmol), 2-[2-phenyl-5-methyl-oxazole-4-yl]-ethanol (Maybridge; 24 mg, 0.116 mmol) and Ph_3P (40 mg, 0.151 mmol) in dry THF (3 mL) was added dropwise DEAD (26 mg, 0.151 mmol). The solution was stirred at RT overnight.
- 10 The orange-red solution was concentrated *in vacuo* and the residue was purified by Prep-HPLC (continuous gradient from 50% A:50% B to 100%B; A = 90% H_2O :10 %MeOH + 0.1% TFA); (B = 90% MeOH/10% H_2O + 0.1% TFA) for 10 min; YMC SH-343-5 ODS 20 x 100 mm (5 μm) column) to provide title
- 15 compound (30 mg, 47%) as a yellow viscous oil.

D.




20

- Part C compound was dissolved in MeOH (3 mL) and H_2O (0.3 mL). To this solution was added LiOH (3 mg) and the reaction was stirred at RT for 3 h. Volatiles were removed *in vacuo* and the solution was acidified with 1N
- 25 aqueous HCl to pH = ~3-4. The aqueous phase was extracted with EtOAc (3 x 10 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated *in vacuo* to give title compound as a white solid (18 mg; 64%). LC/MS (electrospray) gave the
- 30 correct molecular ion $[(\text{M}+\text{H})^+ = 516]$.


5

CC1=C(C(=O)OCCN(C1=O)COC2=CC=C(C=C2)Cl)CCOC3=CC=C(C=C3)OCC4=CC=C(C=C4)N=C5C(=C(C=C5)N)C6=CC=CC=C6

A.



15



20

COC(=O)CN(Cc1ccc(OCC2=C(C)N(C2=O)c3ccccc3)cc1)C(=O)Oc4ccc(Cl)cc4

- 201 -


10

CC1=C(CCOc2ccc(cc2)COC(=O)NCC(=O)O)c3nc(C4=CC=CC=C4)oc31

A solution of Part B compound and LiOH.H₂O (5 mg) in THF:H₂O (4:1) was stirred at RT for 1 h. The reaction solution was acidified to pH 3 with aqueous HCl, then extracted with EtOAc. The combined organic extracts were concentrated in vacuo to give title compound (5 mg; 18%) as a white solid. [M + H]⁺ = 535.2; 537.2


CC1=C(CCOc2ccc(cc2)CN(CCC(=O)O)C(=O)Oc3ccc(Cl)cc3)C(=O)c4ccccc4N1




Title compound was synthesized using the same sequence as in Example 465 with the exception that the aldehyde




Following procedures as described above, Examples 467 to 472 compounds were prepared.

10



Example No.	R ³	[M+H] ⁺
467		501.3
468		563.3
469		515.3


Examples 470 to 472

5


Example No.	R ³	[M+H] ⁺
470		501.3
471		563.3
472		515.3

Example 473

10



A.



15

A mixture of 3-iodophenol (2.0 g; 9.1 mmol), acetic anhydride (4.6 g; 45.5 mmol) and pyridine (3.6 g; 45.5 mmol) was stirred in CH₂Cl₂ (20 mL) for 3 h. The resulting mixture was washed with saturated aqueous NH₄Cl

20

5

C[Si](C)(C)C#Cc1ccc(O)cc1

15

CC(=O)Oc1ccccc1C#CSi(C)(C)C


To a solution of crude Part B compound in CH_2Cl_2 (2 mL) were added pyridine (3 mL; 37 mmol) and acetic anhydride (4 mL; 42 mmol). The reaction was stirred at RT for 2 h, then was partitioned between saturated aqueous NH_4Cl (30 mL) and CH_2Cl_2 . The organic phase was washed with additional saturated aqueous NH_4Cl (30 mL) and H_2O (100 mL), dried (Na_2SO_4) and concentrated in vacuo to give Part C compound, which was used in the next step without further purification.

30

CC(=O)Oc1ccc(C#C)cc1CC(=O)Oc1ccccc1C=C

20


F.



30


5

10




15

25




30

CC1=C(CCOc2ccc(cc2)C(=O)N(C)C(=O)O)N(C1)c3ccccc3

A solution of Part H compound and LiOH.H₂O in MeOH/H₂O (10 mL of a 9:1 mixture) was stirred at RT overnight. The solution was then acidified to pH ~3 with aqueous HCl and extracted with EtOAc. The combined organic extracts were concentrated in vacuo and purified by preparative HPLC to give title compound (10.1 mg; 95%) as an off-white lyophilate. [M + H]⁺ = 527.3


20

CCOC(=O)COc1ccccc1Oc2ccccc2

- 208 -

A mixture of 3-benzyloxybenzaldehyde (2.00 g; 1.0 mmol), ethyl bromoacetate (1.67 g; 1.0 mmol) and Cs_2CO_3 (3.25 g; 1.0 mmol) in DMF (20 mL) was stirred at RT for 8 h. The reaction mixture was partitioned between H_2O (300 mL) and EtOAc (100 mL). The aqueous phase was extracted with EtOAc (2 x 100 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; 85:15 hex:EtOAc) to obtain Part A compound (3.48 g; >100%) as a colorless oil.


B.



15

To a solution of Part A compound (3.4 g; 11.9 mmol) in dry THF (50 mL) under Ar was added LiAlH_4 (36 mL of a 0.5 M solution in THF; 17.8 mmol) dropwise. The reaction was stirred at RT for 1 h. The reaction was quenched by slow addition of saturated aqueous NH_4Cl (1 mL). Volatiles were removed in vacuo and the residue was partitioned between EtOAc (100 mL) and 1 M aqueous HCl. The organic phase was dried (Na_2SO_4) and concentrated in vacuo to give Part B compound (2.4 g; 98%) as a white solid.

C.




30

To a solution of Part B compound (2.4 g; 9.8 mmol) and Ph_3P (3.1 g; 14.7 mmol) in CH_2Cl_2 was added CBr_4 (4.80 g; 14.7 mmol). The reaction was stirred at RT overnight, then concentrated in vacuo. The residue was

chromatographed (SiO_2 ; 95:5 hex:EtOAc) to give Part C compound (2.8 g; 93%) as a white solid.

D.




5

A mixture of Part C compound (310 mg; 1.0 mmol) and potassium tert-butoxide (113 mg; 2.0 mmol) in toluene (20 mL) was heated at 105°C for 20 min. Additional K₂CO₃ (56 mg; 1.0 mmol) was added and the reaction heated at 105°C for another 10 min. The mixture was cooled to RT and partitioned between H₂O (100 mL) and EtOAc (100 mL). The organic phase was washed with H₂O (2 x 100 mL), dried (Na₂SO₄) and concentrated in vacuo. The reaction was repeated with additional Part C compound (500 mg; 1.63 mmol) and K₂CO₃ (182 mg; 1.63 mmol). The combined crude reaction products were chromatographed (SiO_2 ; hexane) to give Part D compound (590 mg; 89%) as a colorless oil.

20


E.



To a 0°C solution of Part D compound (1.4 g; 6.2 mmol) in DCE (100 mL) was added neat diethylzinc (1.6 mL; 16 mmol) dropwise, followed by ICH₂Cl (5.46 g; 31 mmol). The reaction mixture was allowed to warm to RT and stirred at RT overnight, then washed with 1M aqueous HCl. The organic phase was dried (Na₂SO₄) and concentrated in vacuo. The crude residue was chromatographed twice (SiO_2 ; hexane) to give Part E compound (510 mg; 30%) in addition to recovered starting material Part D compound (250 mg; 18%).

30

F.




5

To a -78°C solution of Part E compound (510 mg; 2.2 mmol) in liquid NH_3 (30 mL) was added Na (500 mg; 22 mmol). The dark blue solution was stirred at -78°C for 4 h, then was allowed to warm to RT overnight. The remaining solid residue was partitioned between 1 M aqueous HCl and EtOAc (50 mL each). The organic phase was dried (Na_2SO_4) and concentrated in vacuo. The crude product was chromatographed (SiO_2 ; 9:1 hexane:EtOAc) to give Part F compound (240 mg; 75%) as a yellow oil.


15

G.



20

To a solution of Part F compound (150 mg; 1.0 mmol) in DMF (10 mL) were successively added K_{Ot}Bu (112 mg; 1.0 mmol) and a solution of the following carbamoyl chloride (44 mg; 1.0 mmol) in DMF (0.5 mL).



25

(prepared as described in Examples 5 and 139). The reaction was stirred at RT for 15 min, after which analytical HPLC indicated that all starting material had been consumed. The mixture was partitioned between H_2O and EtOAc (100 mL each). The organic phase was washed with H_2O (2 x 100 mL), dried (Na_2SO_4) and concentrated in vacuo. The crude product was

30


H.



20

CC1=C(C(=N1)C2=CC=CC=C2)CCOCC3=CC=C(C=C3)COC(=O)NCC(=O)Oc4ccc(OC5C6CC6)cc4C(=O)O


Title compound was synthesized from Example 474 Part F compound (150 mg; 1.0 mmol) and the carbamoyl chloride (440 mg; 1.0 mmol)



(prepared as described in Examples 6 and 139) followed by LiOH/H₂O hydrolysis in the same manner as in Example 474.

- 5 The title compound was isolated and purified as a colorless oil (340 mg; 92% over 2 steps). [M + H]⁺ = 543.3

Following the procedures described above, the Examples 476 to 494 compounds were prepared.




5

Example No.	R ^{3h}	[M+H] ⁺
476		519.1
477		535.1
478		579.1; 581.0
479		535.3
480		551.3
481		595.3; 597.3
482		529.3
483		527.3
484		543.4

CC1=C(C(=O)O)N(Cc2ccc(OCC3C(=N1)C(=C3)C4=CC=CC=C4)cc2)C(=O)OR3h

- 215 -

Example 492




5

Example 492 was synthesized according to the procedures described hereinbefore.


¹H NMR (CDCl₃; 400 MHz): δ 0.68 (t, J = 4.4 Hz; 2H), 0.94 (t, J = 4.4 Hz; 2H), 1.87 (m, 1H), 2.42 (s, 3H), 3.06 (s, 2H), 4.02 (t, J = 5.2 Hz, 2H), 4.22 (t, J = 5.2 Hz, 2H), 4.60 (2 peaks, 2H), 6.84-6.89 (m, 4H), 7.15-7.26 (m, 4H), 7.40-7.47 (m, 3H), 7.98-8.00 (m, 2H).

15 The required (commercially unavailable) phenols and chloroformates for the synthesis of the above carbamate-acid analogs were prepared as follows:

3-Fluoro-4-methyl-phenyl chloroformate




A.



25 A mixture of 5-methoxy-2-methyl aniline (5.0 g; 36 mmol), HCl (7.6 mL of a 12 M solution; 91 mmol) and H₂O (11 mL) was heated at 60°C for 15 min until complete dissolution had occurred. The reaction was cooled to 0°C


and an aqueous solution of NaNO_2 (2.5 g; 36 mmol) was added dropwise (internal temperature $\leq 7^\circ\text{C}$). The reaction was stirred at 0°C for 30 min and a 0°C solution of HBF_4 (5.3 mL of a 48% solution; 40 mmol) was added cautiously. 5 The reaction was stirred at 0°C for 20 min, and the resultant brown solid was filtered, washed with ice water (3 x 10 mL) and H_2O (2 x 10 mL). The solid was dried under high vacuum for 20 h, then heated (heat gun) until evolution of BF_3 (white fumes) had ceased. The resulting 10 brown oil was partitioned between EtOAc and H_2O . The organic phase was dried (Na_2SO_4), concentrated in vacuo and distilled by Kugelrohr to give 3-fluoro-4-methyl anisole (1.6 g; 31%) as a colorless oil.

15 B.



To a -70°C solution of 3-fluoro-4-methyl anisole (1.62 g; 11.6 mmol) in CH_2Cl_2 (10 mL) was added dropwise 20 BBr_3 (10 mL; 12 mmol). The reaction mixture was stirred at -70°C for 10 min, then allowed to warm to 0°C and stirred at 0°C for 2 h. The reaction was allowed to warm to RT and concentrated in vacuo and the residue was partitioned between H_2O and EtOAc . The organic phase was 25 washed with H_2O , dried (Na_2SO_4) and concentrated in vacuo to give 3-fluoro-4-methyl phenol (1.1 g; 75%) as an oil.

C.




30

A solution of 3-fluoro-4-methyl phenol (1.1 g; 8.7 mmol), phosgene (5.9 mL of a 1.93 M solution in toluene;


5

10



15

20



25


(1.72 g; 34%) as an oil. This phenol was converted to 3-bromo-4-methyl phenyl chloroformate (1.9 g; 82%) using the same procedure (phosgene/ dimethylaniline/heat) as for 3-fluoro-4-methyl phenyl chloroformate above.

5

2-Methoxyphenyl chloroformate (1.5 g) and 3-methoxyphenyl chloroformate (1.5 g) were both synthesized in the same way as for 3-fluoro-4-methyl phenyl chloroformate (phosgene/dimethylaniline/heat) from 2-methoxyphenol (2 g) and 3-methoxyphenol (2 g) respectively.

10

3-chloro-4-methoxy phenol




15

To a 0°C solution of 3-chloro-4-methoxy aniline (1.0 g; 6.4 mmol) in 1:1 H₂O:conc. H₂SO₄ (100 mL) was added very slowly a solution of NaNO₂ (0.5 g; 7.6 mmol) in H₂O (10 mL). Thick yellow fumes were emitted, and the black solution was then heated to reflux for 30 min. The mixture was extracted with EtOAc (4 x 50 mL) and the combined extracts were concentrated in vacuo. The residue was chromatographed (SiO₂; 4:1 hex:EtOAc) to obtain 3-chloro-4-methoxy phenol (300 mg; 30%) as a yellow oil.

20

25

3-Fluoro-4-methoxy-phenol




30

A solution of 3'-fluoro-4'-methoxyacetophenone (10 g; 59 mmol) and m-chloroperbenzoic acid (50% purity; 30


g; 89 mmol) in CH_2Cl_2 (300 mL) was stirred at RT overnight. The solution was washed with saturated aqueous Na_2CO_3 , then filtered through a pad of SiO_2 (CH_2Cl_2 as eluent) and finally chromatographed (SiO_2 ; hex:EtOAc 4:1) to give the crude product (3'-fluoro-4'-methoxy phenyl acetate; 63 g). A solution of this crude material and $\text{LiOH}\cdot\text{H}_2\text{O}$ (5 g; 120 mmol) in $\text{MeOH}:\text{H}_2\text{O}$ (100 mL of a 9:1 mixture) was stirred at RT overnight. Volatiles were removed in vacuo, and the residue was partitioned between excess aqueous 1 M HCl and EtOAc (aqueous layer pH~3). The aqueous phase was extracted with EtOAc (2x). The combined organic extracts were dried (Na_2SO_4) and concentrated in vacuo to give 3-fluoro-4-methoxy phenol (6.1 g; 72%) as an oil.

3-bromo-4-methoxy phenol (4.39 g; 47% for 2 steps) was synthesized using the exact analogous sequence starting from 3-bromo-4-methoxy benzaldehyde.

3-propyl phenol




A.



A mixture of 3-iodoanisole (2 g; 8.5 mmol), trimethylsilylacetylene (1.67 g; 17 mmol), CuI (32 mg; 0.17 mmol) and $(\text{Ph}_3\text{P})_2\text{PdCl}_2$ (59 mg; 0.085 mmol) in diethylamine (10 mL) was stirred at RT for 1 h. Volatiles were removed in vacuo, and the residue was partitioned between EtOAc and brine. The organic phase was washed with brine (2 x 10 mL) and then filtered through a pad of SiO_2 . Volatiles were removed in vacuo to give the crude product (3-trimethylsilylethynyl anisole)

as a light yellow oil. A solution of this crude product and tetrabutylammonium fluoride (6.6 g; 26 mmol) in THF (10 mL) was stirred at RT for 15 min. Volatiles were removed in vacuo and the residue was chromatographed (SiO₂; 9:1 hex:EtOAc) to furnish Part A compound (1.0 g; 89%) as a yellow oil.

B.




10

To a 0°C solution of Part A compound (1.0 g; 7.6 mmol) in anhydrous THF (5 mL) was added dropwise n-BuLi (4.5 mL of a 2.0 M solution in hexane; 9.1 mmol). The resulting yellow solution was stirred at 0°C for 30 min. Methyl iodide (1.6 g; 11.4 mmol) was then added and the reaction was allowed to warm to RT and stirred at RT for 30 min. Volatiles were removed in vacuo and the residue was partitioned between aqueous 1N HCl and EtOAc. The aqueous phase was extracted with EtOAc (3 x 20 mL), and the combined organic extracts were dried (MgSO₄) and concentrated in vacuo to give Part B compound (1.0 g; 92%) as a yellow oil.

15

20

C.




25

A solution of Part B compound (1.0 g) in MeOH (5 mL) was stirred over 10% Pd/C (10 mg) under an atmosphere of H₂ overnight. The catalyst was removed by filtration through a pad of Celite® and the filtrate was concentrated in vacuo to give Part C compound (1.0 g; 100%) as a yellow oil.

30


CCCC1=CC=CC=C1O

Example 495

NC(=O)Cc1ccccc1


- 222 -

B.



5


To a 0°C solution of Part A compound (49 mg; 0.22 mmol) and the aldehyde (50 mg; 0.10 mmol)



- 10 in DCE (3 mL) was added NaBH(OAc)₃ (30 mg; 0.42 mmol). The reaction was allowed to warm to RT and stirred at RT for 2 h, then at 60°C for 18 h. The mixture was cooled to RT and concentrated in vacuo. The residue was purified by preparative HPLC (YMC S5 ODS 30 x 250 mm
- 15 column; flow rate = 25 mL/min; 20 min continuous gradient from 70:30 B:A to 100% B, where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B = 90:10:0.1 MeOH:H₂O:TFA) to give Part B compound.

20


C.




- A solution of crude Part B compound in THF (1 mL) and aqueous LiOH (0.3 mL of a 1 M solution; 0.3 mmol) was stirred at RT for 3 h, then acidified to pH ~3 with aqueous 1 M HCl. The aqueous phase was extracted with EtOAc (2x); the combined organic extracts were concentrated in vacuo. The residue was purified by
- 30 preparative HPLC (YMC S5 ODS 30 x 250 mm column; flow rate = 25 mL/min; 22 min continuous gradient from 70:30 B:A to

$$[M + H]^+ = 486.3$$

Example 496



A.

CC1=C(CCOc2ccc(C=O)cc2)N(=C1)c3ccccc3

in MeOH (2 mL) was added portionwise NaBH_4 (24 mg; 0.65 mmol), after which the reaction was allowed warm to RT and stirred at RT for 1 h. Volatiles were removed in vacuo and the residue was partitioned between H_2O and EtOAc. The organic phase was dried (Na_2SO_4) and concentrated in vacuo to give the intermediate alcohol as an oil. A solution of the alcohol in CH_2Cl_2 (2 mL) and PBr_3 (1 mL of a 1M solution in CH_2Cl_2) was stirred at RT for 30 min. Volatiles were removed in vacuo and the residue was partitioned between aqueous saturated NaHCO_3 and EtOAc. The organic phase was washed with H_2O , dried (Na_2SO_4) and concentrated in vacuo to give Part A compound (150 mg; 62%) as an oil.


- 224 -

CC1=C(COCC2=CC=C(C=C2)OCC3=CC=CC=C3NC(=O)C4=CC=CC=C4C(=O)O)N=C(C1)C5=CC=CC=C5


A solution of Part A compound (42 mg; 0.11 mmol),
5 Example 500 Part A compound (25 mg; 0.11 mmol) and K_2CO_3
(100 mg; 0.71 mmol) in DMF (1 mL) was stirred at RT for 3
days. The reaction mixture was partitioned between EtOAc
and H_2O . The organic phase was washed with H_2O (2x) and
concentrated in vacuo. The residual oil was dissolved in
10 THF (1 mL) and aqueous LiOH (0.3 mL of a 1 M solution)
was added. The reaction was stirred at RT for 3 h, then
acidified to pH ~3 with aqueous 1 M HCl. The aqueous
phase was extracted with EtOAc (2x); the combined organic
extracts were concentrated in vacuo. The residue was
15 purified by preparative HPLC (YMC S5 ODS 30 x 250 mm
column; flow rate = 25 mL/min; 20 min continuous gradient
from 70:30 B:A to 100% B, where solvent A = 90:10:0.1
 H_2O :MeOH:TFA and solvent B = 90:10:0.1 MeOH: H_2O :TFA) to
give title compound (15 mg; 27% yield over 2 steps) as a
20 white solid. $[M + H]^+ = 486.4$

CC1=CC=C(C=C1)OC(=O)N(C)Cc2ccc(OCCc3c(C)c(Oc4ccccc4)n3)cc2

A.

CC1=C(CCCOS(=O)(=O)C)C(=C2C(=C1)N=C(C2)C3=CC=CC=C3)C

B.




To a solution of Part A compound (850 mg, 2.65 mmol) in DCE (15 mL) were successively added glycine methyl ester hydrochloride (333 mg, 2.65 mmol), Et₃N (554 μ L, 4.0 mmol), NaBH(OAc)₃ (786 mg; 3.7 mmol) and acetic

acid (152 μ L; 2.65 mmol). The reaction mixture was stirred at RT for 6 days, then partitioned between aqueous 1 M NaOH and CH_2Cl_2 . The aqueous phase was washed with H_2O , then concentrated in vacuo. The residue was
5 partitioned between EtOAc and 1 M aqueous HCl. The organic phase was washed with brine, dried (MgSO_4) and concentrated in vacuo to give recovered starting material (Part A compound). The pH of the aqueous HCl phase was
10 adjusted to 10 with excess solid NaOH. This aqueous phase was extracted with EtOAc (60 mL). The organic extract was washed with brine (60 mL), dried (MgSO_4) and concentrated in vacuo to give the crude Part B compound (400 mg; 39%) as an oil, which was used in the next step without further purification.


15

C.




20 To a solution of Part B compound (29 mg; 0.074 mmol) in pyridine (1.0 mL) were added 4-tolyl chloroformate (14 μ L; 0.089 mmol) and DMAP (10 mg). The solution was heated at 61°C for 2h, then cooled to RT and concentrated in vacuo to give crude Part C compound (36
25 mg) as a syrup.

D.




5 Volatiles were removed in vacuo and the residue was
acidified to pH 2 with aqueous 1 M HCl, then extracted
with EtOAc (3 x 40 mL). The combined organic extracts
were dried (Na₂SO₄) and concentrated in vacuo. The crude
product was purified by preparative HPLC (YMC S5 ODS 50 x
10 250 mm column; flow rate = 25 mL/min; continuous 20 min
gradient from 70:30 B:A to 100% B, where A = 90:10:0.1
H₂O:MeOH:TFA and B = 90:10:0.1 MeOH:H₂O:TFA) to give title
compound (28 mg; 72% over 2 steps) as a white solid.
[M + H]⁺ = 515.3

15




20 A.



25


B.



5 To a 0°C solution of the crude Part A compound from above in CH₂Cl₂ (198 mL) was slowly added dropwise BBr₃ (12.0 mL; 127 mmol). The reaction was stirred at 0°C for 3 h, then poured cautiously into a 0°C mixture of saturated aqueous NH₄Cl and EtOAc. The aqueous phase was
10 neutralized by slow addition of solid NaHCO₃, then extracted with EtOAc (2x). The combined organic extracts were washed with brine, dried (MgSO₄) and concentrated in vacuo to furnish Part B compound (7.29 g; 44% over 2 steps).


15

C.




20 To a solution of Part B compound (6.13 g; 29.3 mmol) in dioxane:H₂O (98 mL of a 1:1 solution) were successively added NaHCO₃ (3.2 g; 38 mmol) and 4-methoxyphenyl chloroformate (3.92 mL; 26.4 mmol). The reaction was stirred at RT for 2 h, then partitioned between EtOAc
25 and H₂O. The organic phase was washed with brine, dried (MgSO₄), and concentrated in vacuo to give crude Part C compound (10.0 g; 95%).

D.




- 5 To a solution of Part C compound in MeCN (59 mL) were successively added K_2CO_3 (2.43 g; 17.6 mmol) and the mesylate (4.93 g; 17.6 mmol).



- 10 The reaction was heated at 90°C for 20 h, then cooled to RT. The mixture was partitioned between EtOAc and H_2O . The organic phase was washed with brine, dried ($MgSO_4$) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; stepwise gradient from 8:1 to 3:1 to 1:1
15 hexane:EtOAc) to give Part D compound (3.4 g; 36%).

E.



20

- To a solution of Part D compound (3.4 g; 6.25 mmol) in THF: H_2O (31 mL of a 2:1 solution) was added $LiOH \cdot H_2O$ (0.525 g; 125 mmol). The reaction was stirred at RT overnight for 14 h. EtOAc was added and the solution was
25 acidified with 1 N HCl solution to pH ~2. The organic phase was washed with brine, dried ($MgSO_4$) and concentrated in vacuo. The residue was purified by preparative HPLC (YMC S5 ODS 30 x 250 mm column; flow

10


CC1=C(C(=N1)C2=CC=CC=C2)CCOC3=CC=C(C=C3)C[C@H](C)N(C(=O)OC4=CC=C(C=C4)OC)C(=O)O

20 ¹H NMR (DMSO-d₆; 400 MHz): δ 1.50 (2d, J = 7.0 Hz; 3H),
2.37 (s, 3H), 2.94 (t, J = 6.6 Hz, 2H), 3.74 (s, 3H),
3.84 (m; 2H), 4.21 (t, J = 6.6 Hz, 2H), 5.35 (m, 1H),
6.93 (m, 6H), 7.29 (m, 2H), 7.50 (m, 3H), 7.91 (m, 2H)

CC1=C(C(=O)O)N(C)C(=O)Oc2ccc(OC)cc2C1Cc3ccc(OCCc4c(C)c(Oc5ccccc5)n4)cc3

- 231 -


A.



To a RT mixture of (S)-1-(4-methoxyphenyl)-
5 ethylamine (5.45 g, 36 mmol) in THF (50 mL) and aqueous
NaHCO₃ (6.05g in 25 mL H₂O) was added dropwise benzyl
chloroformate (6.20 mL; 43 mmol). The reaction was
stirred at RT for 30 min; the organic phase was isolated
and concentrated in vacuo. The residue was partitioned
10 between EtOAc and H₂O (100 mL each); the organic phase was
washed with brine, dried (MgSO₄), and concentrated in
vacuo to about 30 mL volume. An equivalent volume of
hexane (30 mL) was added and Part A compound (9.12 g;
89%) crystallized as colorless needles.


15

B.




To a -78°C solution of Part A compound (2.50 g; 8.8
20 mmol) in anhydrous CH₂Cl₂ (11 mL) was added dropwise a
solution of BBr₃ in CH₂Cl₂ (11.4 mL of a 1.0 M solution;
11.4 mmol) over 25 min. The reaction was allowed to warm
to 0°C and stirred at 0°C for 6 h, then quenched carefully
at -78°C by dropwise addition of excess MeOH (6 mL). The
25 solution was allowed to warm to 0°C and stirred at 0°C for
5 min. The solution was partitioned between CH₂Cl₂ (60
mL) and H₂O (50 mL). The organic phase was washed
successively with brine and 5% aqueous NaHCO₃ (50 mL
each), dried (MgSO₄) and concentrated in vacuo. The
30 residue was chromatographed (SiO₂; stepwise gradient from
4:1 to 1:1 hex:EtOAc) to furnish Part B compound (1.30 g;
63% yield based on 650 mg (26%) of recovered unreacted
Part A compound) as a white solid.

C.




A mixture of Part B compound (1.10 g; 4.1 mmol),
5 K_2CO_3 (680 mg; 4.9 mmol) and the mesylate (1.25 g; 4.4
mmol)




10 in MeCN (30 mL) was heated at 90°C for 18 h, then cooled to RT. Volatiles were removed in vacuo, and the residue was partitioned between EtOAc and H₂O (100 mL each). The organic phase was washed with brine (100 mL), dried (MgSO₄) and concentrated in vacuo. The residue was
15 chromatographed (SiO₂; stepwise gradient from 9:1 to 3:2 hexane:EtOAc) to give Part C compound (1.40 g; 78%) as a white solid.

D.




A mixture of Part C compound (1.30 g; 2.85 mmol) and 10% palladium on carbon (200 mg) in MeOH (50 mL) was stirred under an atmosphere of H₂ (balloon) at RT for 2 h, at which point the reaction was complete by HPLC. The catalyst was filtered off through Celite® and the filtrate was concentrated in vacuo to give Part D compound (600 mg; 65%) as a white solid.

E.



5

F.



15

20

25


CC1=C(C(=O)N(C1)C(=O)Oc2ccc(OC)cc2)Cc3ccc(OCCc4cnc(C)c4c5ccccc5)cc3

Example 499A


CC1=C(C(=O)O)N(C)C(=O)Oc2ccc(OC)cc2Oc3ccc(OCCc4c(C)c(Oc5ccccc5)n4)cc3

- 235 -

Alternative Synthesis of Example 498 Part B Compound




5 A.



A mixture of tert-butyldimethylsilyl chloride (357 mg; 2.36 mmol), (alternative) Example 498A Part B compound from above (535 mg; 1.97 mmol) and imidazole (161 mg; 2.36 mmol) in DMF (5 mL) was stirred at RT for 2 h. The reaction was partitioned between EtOAc (20 mL) and water (50 mL). The organic phase was washed with water (2 x 50 mL), dried (Na₂SO₄), and concentrated in vacuo. The residue was chromatographed (SiO₂; hex:EtOAc 3:1) to give Part A compound (320 mg; 42%) as an oil in addition to recovered starting phenol (150 mg; 20%).


20 B.



A mixture of Part A compound (320 mg; 0.83 mmol) and 10% palladium on carbon (30 mg) in MeOH (30 mL) was stirred under an atmosphere of H₂ (balloon) at RT for 1 h, at which point the reaction was complete by HPLC. The catalyst was filtered off through Celite[®] and the filtrate was concentrated in vacuo to give Part B compound (230 mg) as a white solid which was used in the next step without further purification.

30


C.



A solution of Part B compound (230 mg), methyl bromoacetate (86 μ L; 0.91 mmol) and Et₃N (127 μ L; 0.91 mmol) in THF (10 mL) was stirred at RT for 15h. The reaction mixture was partitioned between H₂O and EtOAc (30 mL) each. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo. The residue was chromatographed (SiO₂; stepwise gradient from hex:EtOAc 9:1 to 1:1) to furnish Part C compound (177 mg; 66% over 2 steps) as an oil.

10


Example 498 Part B Compound



To a solution of Part C compound (177 mg; 0.55 mmol) in THF (5.5 mL) was slowly added tetrabutylammonium fluoride (1.65 mL of a 1 M solution in THF). The reaction was stirred at RT for 10 min, then partitioned between water and EtOAc. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo. The residue was purified by preparative HPLC (YMC reverse phase ODS 20 x 100 mm column; flow rate = 20 mL/min; 10 min continuous gradient from 100%A to 100% B + 10 min hold-time at 100% B, where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B = 90:10:0.1 MeOH:H₂O:TFA; retention time = 2.6 min) to provide the title compound (97 mg; 84%).


30

Example 500



CCCC(=O)c1ccc(OCCc2c(C)c3c(c2)nc(c3)C4=CC=CC=C4)cc1


B.



A mixture of Part A compound (3.42 g; 9.42 mmol), glycine methyl ester HCl salt (1.18 g; 9.42 mmol), Et₃N (1.97 mL; 14.1 mmol), NaBH(OAc)₃ (2.80 g; 13.2 mmol) and HOAc (0.54 mL; 9.42 mmol) in DCE (20 mL) was stirred at
25 RT for 6 days. At this point the reaction was incomplete, but was not progressing any further. The reaction was quenched with saturated aqueous NaHCO₃ (6 mL), then concentrated in vacuo. The residue was partitioned between saturated aqueous NaHCO₃ and EtOAc.
30 The organic phase was washed with saturated aqueous NaHCO₃ and H₂O, then extracted with 1M aqueous HCl (the unreacted starting material remained in the organic phase). The aqueous phase was basified with NaOH, then extracted with EtOAc. The organic phase was washed with H₂O and brine,

dried (MgSO_4) and concentrated in vacuo to give Part B compound (365 mg; 9%) as an oil.

C.




5

To a solution of Part C compound (50 mg; 0.11 mmol) in pyridine (1 mL) was added 4-methoxyphenyl
10 chloroformate (40 μL) and DMAP (5 mg). The reaction mixture was heated at 60°C for 6 h, then was cooled to RT and volatiles were removed in vacuo. The residue was dissolved in THF/MeOH/ H_2O (1 mL of a 2:2:1 mixture) and LiOH (30 mg) was added. The reaction was stirred at RT
15 for 18 h, then was acidified with aqueous 1 M HCl to pH ~ 2. The mixture was extracted with EtOAc (30 mL), washed with H_2O and brine (15 mL each), dried (MgSO_4) and concentrated in vacuo to give the crude product. This material was purified by preparative HPLC (YMC S5 ODS 30
20 x 250 mm column; continuous gradient from 60:40 A:B to 100% B over 30 min) to give, after lyophilization from MeOH/ H_2O , the title compound (52 mg; 79%) as a white solid. $[\text{M} + \text{H}]^+ = 573.3$

25

Example 501




CCOC(=O)CN=Cc1ccc(OCCCC2=C(C)OC(=N2)c3ccccc13)cc1CC1=C(CCOc2ccc(C=O)cc2)N(=C1)c3ccccc3

15

COC(=O)CN(C=C)Cc1ccc(OCCc2c(C)c(Oc3ccccc3)n2)cc1


- 240 -

C.



5 To a 0°C solution of Part B compound (150 mg; 0.36 mmol) and Et₃N (51 µL; 0.36 mmol) in CH₂Cl₂ (4 mL) was added dropwise 4-methoxyphenyl chloroformate (53 µL; 0.36 mmol). The reaction was allowed to warm to RT and stirred at RT for 1 h, then concentrated in vacuo. The residue was chromatographed (SiO₂; hexane:EtOAc 2:1) to give Part C compound (200 mg; 98%) as an oil.


D.



15


A solution of Part C compound (100 mg, 0.18 mmol) and LiOH·H₂O (30 mg, 0.72 mmol) in THF:MeOH:H₂O (1 mL of a 1:1:1 solution) was stirred at RT for 2 h. The reaction mixture was then acidified to pH ~ 2 with aqueous 1N HCl. The aqueous phase was extracted with EtOAc (2x). The combined organic extracts were dried (Na₂SO₄), concentrated in vacuo and lyophilized from dioxane to provide title compound (80 mg; 82%) as a white solid.

25 [M + H]⁺ = 557.2

Example 502

5


A.



A solution of Example 501 Part C compound (100 mg; 0.18 mmol) in MeOH (10 mL) in the presence of 10% Pd/C (50 mg) was stirred under an H₂ atmosphere for 2 h at RT. The catalyst was then filtered off using a pad of Celite[®]. The filtrate was concentrated in vacuo to give Part A compound (100 mg; 100%) as an oil.


15

B.




20

Title compound (87 mg; 90%; white solid lyophilate) was obtained from Part A compound in the same way as Example 501 compound was synthesized from Example 501 Part C compound. $[M + H]^+ = 559.2$


Example 503

5 A.




To a solution of 5-methyl 2-phenyl-thiazol-4-yl-ethanol (50 mg; 0.23 mmol) in CH_2Cl_2 (3 mL) were successively added Et_3N (50 μL ; 0.36 mmol) and methanesulfonyl chloride (20 μL ; 0.26 mmol). The reaction was stirred at RT for 2 h, then was partitioned between CH_2Cl_2 and aqueous 1 M HCl. The organic phase was washed with brine, dried (Na_2SO_4), and concentrated in vacuo to give Part A compound (68 mg; 100%) as a colorless oil. This material was used in the next step without further purification.

20 B.



A mixture of the phenol (prepared using the identical procedures as described for the synthesis of Example 498 Part C compound except that ethyl bromoacetate was used instead of methyl bromoacetate)




10


CC1=C(C(=N1)C2=CC=CC=C2)CCOC3=CC=C(C=C3)C[C@H](C)N(C(=O)OC4=CC=C(C=C4)OC)C(=O)O

20

Example 504




5 The title compound was prepared in exactly the same way as for Example 503 except that the [S]-enantiomer




was used for the alkylation step. $[M + H]^+ = 547.2$

Example 505



15 A.



To a solution of 1-(4-methoxyphenyl)-1-cyclopropane-
20 carboxylic acid (250 mg; 1.3 mmol) in dioxane (8 ml) were
successively added Et₃N (198 μL; 1.43 mmol) and diphenyl-
phosphoryl azide (307 μL; 1.43 mmol). The reaction was
stirred at RT for 5 min, then heated to 80°C for 3 h.

10

COC(=O)CN(C1CC1)c2ccc(OC)cc2


20

COC(=O)CN(C1CC1)c2ccc(O)cc2

To a 0°C solution of Part B compound (90 mg; 0.38 mmol) in CH₂Cl₂ (12.7 mL) was slowly added neat BBr₃ (82 µL; 0.87 mmol). The reaction was stirred at 0°C for 3 h, then was partitioned between ice cold saturated aqueous

NH₄Cl and EtOAc. The organic phase was discarded and the aqueous layer was neutralized by addition of NaHCO₃, then extracted with EtOAc (2x). The combined organic extracts were washed with brine, dried (MgSO₄), and concentrated in vacuo to give Part C compound (50 mg; 59%).


D.



10

A mixture of Part C compound (50 mg; 0.22 mmol), 4-methoxyphenyl chloroformate (33 mg; 0.22 mol) and NaHCO₃ (25 mg; 0.29 mmol) in 1:1 aqueous dioxane (7.5 mL) was stirred at RT for 2 h. The reaction mixture was partitioned between EtOAc and H₂O. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo to give Part D compound (45 mg; 52%).


E.



20

A mixture of Part D compound (45 mg; 0.12 mmol), K₂CO₃ (30 mg; 0.22 mol) and the mesylate (33 mg; 0.12 mmol)

25




5

CC1=C(C2=CC=CC=C2O1)CCOC3=CC=C(C=C3)C4(C5C6C5C(C6)C)N(C4)CC(=O)Oc7ccc(OC)cc7C(=O)O


15


20



Example No.	R ^a	[M+H] ⁺
506	(±) -Me	515.3
507	(±) n-Bu	557.4

5




Example No.	R ^a	[M+H] ⁺
508	(±) Me	531.3
509	(±) Et	545.1
510	(±) i-Bu	573.3
511	(±) 	571.3

CC1=CN(C(=O)C1)CCOC2=CC=C(C=C2)C(C)N(C)C(=O)Oc3ccc(C)cc3C(=O)O

¹H NMR (DMSO-d₆; 400 MHz): δ 1.47 and 1.54 (2d, J = 7.5 Hz; 3H), 2.29 (s, 3H), 2.37 (s, 3H), 2.93 (t, J = 6.6 Hz, 2H), 3.81 (2d, J = 18 Hz; 2H), 4.21 (t, J = 6.6 Hz, 2H), 5.3 (m, 1H), 6.94 (m, 4H), 7.18 (d, J = 8.4 Hz, 2H), 7.31 (m, 2H), 7.49 (m, 2H)


CC1=C(C(=O)OCC1c2ccc(OCC3=C(C)OC(=O)N(C)C(=O)Oc4ccc(OC)cc4)cc2)c5ccccc5

¹H NMR (DMSO-d₆; 400 MHz): δ 1.47 and 1.54 (2d, J = 7.5 Hz; 3H), 2.37 (s, 3H), 2.94 (t, J = 6.6 Hz, 2H), 3.74 (s, 3H), 3.81 (m, 2H), 4.21 (t, J = 6.6 Hz, 2H), 5.36 (m, 1H), 6.94 (m, 4H), 7.29 (m, 2H), 7.49 (m, 3H), 7.91 (m, 2H)






Example No.	Structure	[M+H] ⁺
512	 (±)	531.3


5 The synthesis of Examples 513-518 involved the use of Example 541 Part B compound as the alkylating agent.



Example No.	Structure	[M+H] ⁺
513		517.2
514		517.2
515		501.2


Example No.	Structure	[M+H] ⁺
516		501.2
517		517.2
518		517.2

Example 519




5

A.



10


A mixture of methyl α -aminoisobutyrate hydrochloride (108 mg; 0.7 mmol), Et₃N (146 μ L; 111 mmol), NaBH(OAc)₃ (222 mg; 11 mmol) and the aldehyde (215 mg; 0.7 mmol)



15

in DCE (5 mL) was stirred at RT for 21 h. Some starting material remained, so the reaction was heated at 55°C for 4 h (no further reaction). Saturated aqueous NaHCO₃ was added, and volatiles were removed in vacuo. The residue was partitioned between H₂O and EtOAc. The aqueous phase was extracted with EtOAc (2x). The combined organic extracts were washed with brine and extracted with aqueous 1 M HCl. The aqueous phase was basified with solid NaOH and extracted with EtOAc (2x). The organic extracts were dried (Na₂SO₄) and concentrated in vacuo to give crude Part A compound (174 mg; 61%).


B.



15

A solution of Part A compound (120 mg; 0.29 mmol) aqueous LiOH (2.0 mL of a 0.3 M solution of a 1:1:1 mixture of THF:MeOH:H₂O) was stirred at RT overnight. The reaction was acidified to pH ~ 2 with aqueous 1 M HCl, then was concentrated in vacuo and purified by preparative HPLC (YMC S5 ODS 30 x 250 mm column; flow rate = 25 mL/min; continuous gradient from 40:60 B:A to 100% B over 30 min, where solvent A = 90:10:0.1 H₂O:MeOH:TFA; solvent B = 90:10:0.1 MeOH:H₂O:TFA) to furnish Part B compound (60 mg; 53%) as a syrup.

C.






30


10

15



CC1=C(CCOc2ccc(cc2)CN(R3)(Rb)C(Rc)C(=O)O)N=C(N1)c3ccccc3








The image displays four chemical structures, labeled 1a, 1b, 1c, and 1d, arranged vertically. Each structure consists of a 2-methyl-2-phenyl-1,3-oxazole ring connected via a 2-ethoxyethyl linker to a 4-substituted phenoxy group. The substituent at the para position of the phenoxy group is an ester linkage (-O-C(=O)-N-CH(CH₃)-CO₂H) where the nitrogen is also bonded to a 4-substituted phenyl ring. The substituents on the terminal phenyl rings are: 1a) 4-methoxyphenyl, 1b) 4-methylphenyl, 1c) 4-methoxyphenyl, and 1d) 4-phenylphenyl.

Example No.	Structure	[M+H] ⁺
524		531.2
525		515.2
526		515.2




5

Example No.	Structure	[M+H] ⁺
527		543.3
528		527.3


529		545.3
530		531.2
531		515.2
532		515.2
533		531.2
534		515.2
535		515.2

Example 536



5


A.



10 To a 0°C solution of (R)-(-)-lactate (3.0 g; 29 mmol) and Et₃N (4.8 mL; 35 mmol) in CH₂Cl₂ (60 mL) was added methanesulfonyl chloride (2.67 mL; 35 mmol). The mixture was stirred at 0°C for 1 h, then partitioned between CH₂Cl₂ and 1M aqueous HCl (100 mL each). The organic phase was washed with H₂O and brine, dried (MgSO₄), and concentrated in vacuo without heating to give Part A compound as an oil (4.5 g; 86%), which was used in the next step without further purification.

20


B.



25 A mixture of Part A compound (1.42 g; 6.0 mmol), (R)-4-methoxy-α-methyl benzylamine (300 mg, 2.0 mmol), and K₂CO₃ (828 mg; 6.0 mmol) in MeCN (20 mL) was heated at 70°C for 17 h (some amine starting material still remained). The reaction cooled to RT, filtered, and the volatiles were removed in vacuo. The residue was partitioned between EtOAc and H₂O. The organic phase was washed with brine, dried (MgSO₄), and concentrated in

30

5 C.




20

CC(C)(C(=O)OC1=CC=C(C=C1)OC)N(C(=O)OC2=CC=C(C=C2)OC)C3=CC=C(C=C3)O

30

gradient from 3:1 to 1:1 hexane:EtOAc) to furnish Part D compound (330 mg; 66%).


E.



5

To a solution of Part D compound (330 mg; 0.88 mmol) in MeCN (20 mL) were successively added K_2CO_3 (165 mg; 1.2 mmol) and the mesylate (337 mg; 1.2 mmol).

10




15

The reaction mixture was heated at 95°C for 16 h, then cooled to RT and filtered. The filtrate was concentrated in vacuo and then partitioned between EtOAc and H_2O . The organic phase was washed with brine, dried ($MgSO_4$) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; 3:1 hexane:EtOAc) to give Part E compound (350 mg; 71%).

20

F.



25


To a solution of Part E compound (350 mg; 0.62 mmol) in THF: H_2O (15 mL of a 2:1 solution) was added $LiOH \cdot H_2O$ (52 mg; 1.2 mmol). The reaction was stirred at RT overnight for 14 h; then EtOAc was added and the

10 $[M + H]^+ = 545.3$


Alternative Synthesis of Example 536:




A.




20 prepared from (R)-1-(4-methoxyphenyl)ethylamine as for
the alternative synthesis of Example 498],




25 K_2CO_3 (400 mg; 2.89 mmol) and the mesylate (710 mg; 2.52 mmol)



10 B.



C.





$$\text{H}_3\text{CO}_2\text{SO}-\overset{\text{CH}_3}{\underset{\text{H}}{\text{C}}}-\text{CO}_2\text{CH}_3$$

- 261 -


filtrate was concentrated in vacuo, and the residue was chromatographed (SiO₂; stepwise gradient from hex:EtOAc 3:1 to 1:1) to furnish Part C compound (98 mg; 47%) as an oil. This intermediate was then used for the preparation of Example 536 in an identical manner to that previously shown.

Examples 537 to 539

10 Following the procedures set out hereinbefore, the Examples 537 to 539 compounds were prepared.


Example No.	Structure	[M+H] ⁺
537		545.3
538		545.3
539		545.3

Example 540



5


A.



10 To a 0°C solution of 5-methyl-2-phenyl-oxazol-4-yl
ethanol (5.0 g; 24.6 mmol), acetone cyanohydrin (3.35 mL;
36.9 mmol) and Ph₃P (7.5 g; 29.5 mmol) in THF (60 mL) was
added DEAD (6.0 mL; 36.9 mmol) dropwise. After addition
was complete, the reaction mixture was warmed to RT and
15 stirred overnight at RT. Volatiles were removed in
vacuo, and the residue was chromatographed (SiO₂;
hexane:EtOAc 2:1) to give Part A compound (4.5 g; 86%) as
an oil.


20

B.




25 A solution of Part A compound (4.5 g; 21.2 mmol)
and H₂SO₄ (concentrated; 20 mL) in EtOH (100 mL) was
heated under reflux overnight. The solution was
concentrated in vacuo to 1/3 its original volume, then
EtOAc (150 mL) and H₂O (100 mL) were cautiously added.
The organic phase was washed with saturated aqueous NaHCO₃,
30 (2 x 100 mL) and brine (150 mL), dried (MgSO₄), and
concentrated in vacuo to give a crude oil. This material

C.




D.




- 264 -

E.



A mixture of Part D compound (380 mg; 1.29 mmol),
5 4-hydroxybenzaldehyde (188 mg; 1.55 mmol) and K_2CO_3 (214
mg; 1.55 mg) in MeCN (12 mL) was refluxed in an oil bath
for 17 h. At this point all starting Part D compound had
been consumed (but there was a significant quantity of
the hydrolysis by-product, Part C compound) by HPLC/MS.
10 The reaction was cooled to RT and the solid precipitates
were filtered off. The filtrate was concentrated in
vacuo and partitioned between EtOAc (60 mL) and H_2O (40
mL). The organic phase was washed with brine (40 mL),
dried ($MgSO_4$), and concentrated in vacuo to give the crude
15 product. This material was chromatographed (SiO_2 ;
stepwise gradient from 4:1 to 1:2 hex:EtOAc) to give Part
E compound (150 mg; 36%) as an oil in addition to Part C
compound (100 mg; 36%).


20 F.



A mixture of Part E compound (150 mg; 0.50 mmol),
25 glycine methyl ester hydrochloride (75 mg; 0.60 mmol) and
 Et_3N (84 μL ; 0.60 mmol) in MeOH (5 mL) was stirred at RT
for 6 h, after which $NaBH_4$ (50 mg) was added cautiously
portionwise. The reaction mixture was stirred at RT
overnight, after which volatiles were removed in vacuo.
30 The residue was partitioned between EtOAc and H_2O . The
organic phase was washed with brine, dried (Na_2SO_4) and

concentrated in vacuo to give Part F compound (180 mg; 97%) as an oil.


G.




A mixture of Part F compound (23 mg; 0.060 mmol), Et₃N (10 μ L; 0.66 mmol) and 4-tolyl chloroformate (10 μ L; 0.066 mmol) in CH₂Cl₂ (1 mL) was stirred at RT for 2 h. Volatiles were removed in vacuo and the residue was dissolved in a solution of THF/MeOH/H₂O (1 mL of a 2:2:1 mixture); LiOH.H₂O (14 mg; 0.33 mmol) was added, and the reaction was stirred at RT for 2 h. Volatiles were removed in vacuo, and the residue was partitioned between aqueous 1 M HCl and EtOAc. The organic extract was concentrated in vacuo and the residue was purified by preparative HPLC (YMC ODS S5 30 mm x 250 mm column, continuous 25 minute gradient from 40% B:60% A to 100% B, hold at 100% B for 15 min, where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B = 90:10:0.1 MeOH:H₂O:TFA; flow rate = 25 mL/min) to give the title compound as a white solid (13 mg; 45% over 2 steps). [M + H]⁺ = 515.3

25

Example 541




A.




5 To a solution of benzaldehyde (23.8 g, 234 mmol) in EtOAc (150 mL; pre-saturated with HCl gas) was added 2,3-butanedione mono-oxime (25.0 g, 234 mmol) in one portion and the resulting solution was stirred at RT for 12 h. Analytical HPLC indicated that all starting materials had
10 been consumed. The reaction mixture was concentrated in vacuo to yield Part A compound as a white solid, which was used in the next step without further purification.

B.



To a solution of Part A compound in CHCl_3 (200 mL) was added dropwise POCl_3 (30 mL, 320 mmol). The reaction was stirred for 12 h at 50°C , then was concentrated in vacuo. The brown residue was partitioned between EtOAc (300 mL) and 1N aqueous NaOH. The organic phase was washed with brine, dried, (MgSO_4) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; Et_2O) to give Part B compound (41.5 g; 86%) as a light brown solid (>95% pure by analytical HPLC and ^1H -NMR analysis).

C.



A solution of 4-hydroxybenzaldehyde (20 g, 160 mmol), glycine methyl ester hydrochloride (22 g, 180 mmol) and Et₃N (25 mL, 180 mmol) in MeOH (200 mL) was stirred at RT for 12 h. The reaction mixture was cooled to 0°C and NaBH₄ (9.0 g, 240 mmol) was added portionwise

5

D.



10

15

E.




25

30

EtOAc (250 mL) and brine (100 mL). The aqueous layer was extracted with EtOAc (3 x 100 mL). The combined organic extracts were dried (MgSO_4) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; stepwise gradient
5 from 3:1 to 1:1 hexane:EtOAc) to give Part D compound (23.6 g; 71%) as a colorless oil.

F.




10

A solution of Part D compound (23.6 g, 47.4 mmol) and $\text{LiOH}\cdot\text{H}_2\text{O}$ (4.0 g, 95 mmol) in THF (200 mL) and H_2O (120 mL) was stirred at RT for 4 h. The reaction mixture was
15 then acidified to pH ~ 2 with aqueous 1N HCl. The aqueous phase was extracted with EtOAc (3 x 200 mL). The combined organic extracts were dried (MgSO_4) and concentrated in vacuo to yield an oily residue, which was recrystallized from EtOAc to provide title compound (19.4
20 g; 84%) as a white solid. $[\text{M} + \text{H}]^+ = 487.23$;


^1H NMR (CD_3OD ; 400 MHz): δ 2.32 (s, 3H), 2.46 (s, 3H), 3.99 & 4.04 (2s, 2H), 4.47 & 4.54 (2s, 2H), 5.01 and 5.00 (2s, 2H), 6.99 (d, $J = 8.4$ Hz, 2H), 7.05 (m, 2H), 7.17
25 (d, $J = 8.4$ Hz, 2H), 7.31 (m, 2H); 7.49 (m, 3H), 8.01 (m, 2H);

^1H NMR (CDCl_3 ; 400 MHz): δ 2.31 (s, 3H), 2.44 (s, 3H), 4.00 (s, 2H), 4.55 (2s, 2H), 5.00 (2s, 2H); 6.99 (m, 4H); 7.13 (m, 2H), 7.21 (d, $J = 8.8$ Hz, 2H); 7.31 (m, 2H);
30 7.44 (s, 3H); 8.01 (s, 2H)

Example 542

5


A.



A mixture of 2-phenyl-5-methyl-oxazole-4-acetic
10 acid (470 mg; 2.17 mmol; Maybridge) pyridine N-oxide (830
mg; 8.74 mmol) and acetic anhydride (350 mg; 3.57 mmol)
in toluene (10 mL) was heated at 90°C for 12 h, then
concentrated in vacuo. The residue was then partitioned
between EtOAc and 1M aqueous HCl. The organic phase was
15 washed with saturated aqueous NaHCO₃, brine, dried
(Na₂SO₄) and concentrated in vacuo to give a dark brown
oil. This material was chromatographed (SiO₂; 4:1
hex:EtOAc) to give Part A compound (143 mg; 35%) as an
oil.


20

B.




25 To a 0°C solution of Part A compound (600 mg; 3.21
mmol) and Ph₃P (3.37 g; 12.9 mmol) in CH₂Cl₂ (50 mL) was
added dropwise a solution of CBr₄ (2.13 g; 6.4 mmol) in
CH₂Cl₂ (20 mL). The solution was stirred at 0°C for 2 h,
then allowed to warm to RT and stirred at RT overnight.
30 Volatiles were removed in vacuo and the residue was
chromatographed (85:15 hexane:EtOAc) to furnish Part B
compound (1.08 g; 98%) as a pale yellow solid.

C.




5 To a -78°C solution of Part B compound (1.12 g; 3.26 mmol) in THF (60 mL) was added n-butyllithium dropwise (4.2 mL of a 1.6 M solution in hexane; 6.72 mmol) over 25 min, while maintaining the internal temperature at $\leq -71^{\circ}\text{C}$. The reaction was stirred at -78°C for 1 h, then allowed to warm slowly to 0°C . Paraformaldehyde (305 g) was then added in one portion and the reaction was stirred at 0°C for 3 h and then quenched with saturated aqueous NH_4Cl . The aqueous phase was extracted with EtOAc (2x); the combined organic
10 extracts were washed with brine, dried (Na_2SO_4) and concentrated in vacuo to give a dark oil. This material was chromatographed (SiO_2 ; 3:2 hex:EtOAc) to give Part C compound (466 mg; 67%) as a yellow solid.

20 D.



To a 0°C solution of Part C compound (466 mg; 2.19 mmol) and Et_3N in CH_2Cl_2 was added dropwise methanesulfonyl chloride (190 μL ; 2.45 mmol) and the reaction was stirred at 0°C for 1 h. The mixture was then partitioned between CH_2Cl_2 and cold 1M aqueous HCl . The organic phase was washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The crude product was
30 chromatographed (SiO_2 ; 3:2 hex:EtOAc) to give Part D compound (533 mg; 84%) as an off-white solid.


E.



5 A mixture of Part D compound (198 mg; 0.68 mmol),
4-hydroxybenzaldehyde (96 mg; 0.79 mmol) and K_2CO_3 (141
mg; 1.02 mmol) in CH_3CN (13 mL) was heated at $70^\circ C$ for
3 h, then stirred at RT overnight. Volatiles were
10 removed in vacuo, and the residue was partitioned between
EtOAc and 1 M aqueous NaOH. The organic phase was washed
with brine, dried (Na_2SO_4) and concentrated in vacuo to
give crude Part E compound (190 mg; 88%) as a yellow oil,
which was used in the next step without further
purification.

15


F.



20 A mixture of Part E compound (123 mg; 0.39 mmol),
glycine methyl ester hydrochloride (248 mg; 1.98 mmol)
and Et_3N (600 μL ; 4.3 mmol) in DCE (15 mL) was stirred at
RT for 15 min, after which $NaBH(OAc)_3H$ (262 mg; 1.2 mmol)
was added in one portion. The reaction was stirred for
25 16 h at RT, after which additional $NaBH(OAc)_3H$ (200 mg;
0.94 mmol) was added. Stirring was continued for 3 h,
after which still more $NaBH(OAc)_3H$ (200 mg; 0.94 mmol) was
added. The reaction was stirred at RT for 48 h, after
which all Part E compound had been consumed. The
30 reaction mixture was partitioned between CH_2Cl_2 and
aqueous $NaHCO_3$. The aqueous phase was extracted with
 CH_2Cl_2 (2x). The combined organic extracts were washed
with brine, dried (Na_2SO_4) and concentrated in vacuo. The
crude product was chromatographed (SiO_2 ; stepwise gradient

from 1:1 to 2:3 hex:EtOAc) to give Part F compound (120 mg; 79%) as a colorless oil which solidified on standing.

G.




5

To a solution of Part F compound (180 mg; 0.46 mmol) and pyridine (100 μ L; 1.24 mmol) in CH_2Cl_2 (10 mL) was added 4-methoxyphenyl chloroformate (105 μ L; 0.71 mmol). The reaction was stirred at RT for 3.5 h, then partitioned between aqueous NaHCO_3 and EtOAc. The aqueous phase was extracted with EtOAc (2x). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The crude product was chromatographed (SiO_2 ; hex:EtOAc 3:2) to give Part G compound (232 mg; 93%) as a colorless oil.

15

H.




20

To a solution of Part G compound (232 mg; 0.43 mmol) in THF: H_2O (12 mL of a 5:1 mixture) was added LiOH. H_2O (1.3 mmol). The solution was stirred at RT overnight, then acidified with aqueous 1M HCl and extracted with EtOAc (2x). The combined organic extracts were washed with brine, dried (Na_2SO_4) and concentrated in vacuo. The crude product was purified by preparative HPLC (YMC S5 ODS 30 x 75 mm column, flow rate = 20 mL/min; continuous gradient from 70:30 B:A to 100% B,

30


where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B = 90:10:0.1 MeOH:H₂O:TFA) to give title compound (160 mg; 71%) as a white solid. [M + H]⁺ = 527.2

5

Example 543


A.

10



A solution of 5-methyl-2-phenyloxazole-4-yl-acetic acid (4.0 g; 18 mmol) and concentrated HCl (2 mL) in MeOH (60 mL) was heated at reflux overnight. Volatiles were removed in vacuo; the residue was partitioned between H₂O and EtOAc. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo to give crude Part A compound as a colorless oil (4.00 g; 94%) which was used in the next step without further purification.

B.




25

To a -78°C solution of LDA (15.0 mL of a 2.0 M solution in heptane/THF; 30 mmol; Aldrich) were successively added dropwise a solution of Part A compound (2.3 g; 10 mmol) in THF (6 mL) and HMPA (500 µL; 2.9 mmol). The solution was stirred at -78°C for 30 min, after which methyl iodide (1.87 mL; 30 mmol) was added dropwise. The solution was stirred at -78°C for 1 h,

then was allowed to warm to 0°C and stirred at 0°C for 1 h. The reaction solution was partitioned between saturated aqueous NH₄Cl and EtOAc. The aqueous phase was extracted with EtOAc (2 x 50 mL). The combined organic
5 extracts were dried (MgSO₄) and concentrated in vacuo to give crude Part B compound (1.90 g; 78%) as a colorless oil, which was used in the next step without further purification.


10 C.



To a -78°C solution of LDA (7.0 mL of a 2.0 M in
15 heptane/THF; 14 mmol; Aldrich) were successively added dropwise a solution of Part B compound (1.8 g; 7.3 mmol) in THF (5 mL) and HMPA (500 µL; 2.9 mmol). The solution was stirred at -78°C for 1 h, then a solution of methyl
iodide (1 mL; 11 mmol) was added dropwise. The solution
20 was stirred at -78°C for 1 h, then was allowed to warm to 0°C and stirred at 0°C for 1 h. The reaction solution was then partitioned between saturated aqueous NH₄Cl and EtOAc. The aqueous phase was extracted with EtOAc (2 x 50 mL). The combined organic extracts were dried (MgSO₄)
25 and concentrated in vacuo. The crude product was combined with the product from another reaction (from 670 mg of Part B compound) and chromatographed (SiO₂; 9:1 hexane:EtOAc) to give Part C compound (2.60 g; 95%) as a colorless oil.

30


D.



To a -78°C solution of Part C compound (1.2 g; 4.63 mmol) in THF (3 mL) under an atmosphere of N_2 was added dropwise a solution of LiAlH_4 (1.0 mL of a 1.0 M solution in THF). The reaction was stirred at -78°C for 1 h, then
5 was allowed to warm to 0°C and stirred at 0°C for 30 min. The reaction was quenched by cautious addition of 1M aqueous potassium sodium tartrate followed by H_2O . The aqueous phase was extracted with EtOAc. The combined organic extracts were washed with H_2O , dried (MgSO_4) and
10 concentrated in vacuo to give crude Part D compound (1.01 g; 94%) as an oil, which was used in the next step without further purification.

E.


15



To an 80°C solution of Part D compound (700 mg; 3.0 mmol), Ph_3P (1.2 g; 4.6 mmol) and 4-hydroxybenzaldehyde
20 (406 mg; 3.3 mmol) in THF (10 mL) was added DEAD (720 μL ; mmol) in two portions over 5 min. The solution was stirred at 80°C for 1 h (starting material still remained), then was concentrated in vacuo. The residue was chromatographed (SiO_2 ; stepwise gradient from 9:1 to
25 5:1 hexane:EtOAc) to give Part E compound (160 mg; 16%).

F.


30



A solution of Part E compound (250 mg; 0.75 mmol), glycine methyl ester hydrochloride (141 mg; 1.13 mmol) and Et_3N (157 μL ; 1.13 mmol) in MeOH (30 mL) was stirred at RT overnight. Excess solid NaBH_4 was added cautiously;
35 the reaction was stirred at RT for 1 h, then concentrated

in vacuo. The residue was partitioned between H₂O and EtOAc. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo to give crude Part F compound (300 mg; 98%) which was used in the next
5 reaction without further purification.


G.



10

To a 0°C solution of Part F compound (150 mg; 0.37 mmol) and Et₃N (51 µL; 0.37 mmol) in CH₂Cl₂ (5 mL) was added 4-methoxyphenyl chloroformate (55 µL; 0.37 mmol). The reaction was allowed to warm to RT and stirred at RT
15 for 2 h. Volatiles were removed in vacuo and the residue was chromatographed (SiO₂; 5:1 hexane:EtOAc) to furnish Part G compound (130 mg; 63%).

H.




20

A solution of Part G compound (130 mg) and LiOH.H₂O (39 mg) in H₂O/THF/MeOH (2 mL of a 1:2:2 mixture) was stirred at RT for 2 h. Volatiles were removed in vacuo,
25 and the residue was acidified with 1.0 M aqueous HCl, then extracted with EtOAc. The organic phase was dried (Na₂SO₄) and concentrated in vacuo to give a residue, which was purified by preparative HPLC (YMC S5 ODS
30 reverse phase C18, 30 x 250 mm column; flow rate = 25

5 lyophilate. $[M + H]^+ = 545.4$


Example 544




Title compound was prepared in analogous fashion to Example 543 except that 3-hydroxybenzaldehyde was used instead of 4-hydroxybenzaldehyde (in the preparation of Example 543 Part E compound). $[M + H]^+ = 545.4$

Example 543 Part E compound). $[M + H]^+ = 545.4$


Example 545



A.




A mixture of the acetylene (38 mg; 0.065 mmol)



(synthesized in a completely analogous fashion to Example
5 542 Part G compound with glycine tert-butyl ester
hydrochloride instead of glycine methyl ester
hydrochloride), quinoline (80 mg; 0.62 mmol) and
Lindlar's catalyst (8 mg; Pd/CaCO₃; Aldrich) in MeOH (8
mL) was stirred under an atmosphere of H₂ at 0°C for 20
10 min. Additional Lindlar's catalyst (8 mg; Pd/CaCO₃;
Aldrich) was then added and stirring was continued under
an atmosphere of H₂ at 0°C for 25 min, after which
reaction was complete. The mixture was filtered, and the
filtrate was concentrated in vacuo. The residue was
15 purified by preparative HPLC (YMC S5 ODS 20 x 100 mm
column; flow rate = 20 mL/min; continuous 20 min gradient
from 80:20 B:A to 100% B, where A = 90:10:0.1 H₂O:MeOH:TFA
and B = 90:10:0.1 MeOH:H₂O:TFA) to give Part A compound
(22 mg; 56%) as a colorless oil.

20

B.




25 To a solution of Part A compound (3 mg; 0.005 mmol)
in CH₂Cl₂ (0.5 mL) was added dropwise TFA (0.25 mL) and
the reaction was stirred for 2 h at RT. Volatiles were
removed in vacuo; the residue was dissolved in CDCl₃,

filtered through a cotton plug and concentrated in vacuo to give the title compound (1.5 mg; 55%) as a colorless oil. $[M + Na]^+ = 551.0$

- 5 Following procedures set out above, Examples 546 to 556 compounds were prepared.


Examples 546 to 556



10

Example No.	Structure	$[M+H]^+$
546		515.4
547		531.3
548		503.3
549		529.4
550		531.2


Example No.	Structure	[M+H] ⁺
551		515.2




Example No.	Structure	[M+H] ⁺
552		531.3
553		487.3
554		503.3
555		527.2

CC1=C(C(=O)O)C(=C(C#CCOC2=CC=C(C=C2)CN(C)C(=O)OC3=CC=C(C=C3)C)C=C4C(=C(C=C(C=C4)O)C5=CC=CC=C5)C6=CC=CC=C6)O1


Example 555



A.



10 A mixture of the mesylate (124 mg; 0.43 mmol),




3-hydroxybenzaldehyde (62 mg; 0.51 mmol) and K_2CO_3 (94 mg; 0.68 mmol) in CH_3CN (10 mL) were heated at 70°C for 48 h.

15 The reaction was cooled to RT, EtOAc was added, and the
mixture was washed with aq 1M NaOH and brine. The
organic phase was dried (Na_2SO_4) and concentrated in
vacuo. The residue was chromatographed (SiO_2 ; hex:EtOAc
4:1) to give Part A compound (71 mg; 52%) as a colorless
20 oil. $[\text{M} + \text{H}]^+ = 318.2$

COC(=O)NCc1ccc(OCC#Cc2c(C)c(O)c(Nc3ccccc3)c2)cc1

20 C.



- 283 -

5

Cc1c(C#CCOC2=CC=C(CN(C2)COC(=O)C)C3=CC=C(OC)C3)c4ccccc4n1

20


CC1=C(C(=C(C=C1)C#CCOC2=CC=C(C=C2)CN(C)C(=O)O)OC(=O)C3=CC=C(C=C3)C)C4=CC=CC=C4N=C5C=C(C)OC(=C5)C6=CC=CC=C6

Title compound was synthesized using the same sequence as Example 555 compound from Example 555 Part B compound. Acylation with 4-methyl chloroformate (67% after HPLC purification) followed by LiOH hydrolysis


furnished title compound (5 mg; 57% after HPLC purification). $[M + H]^+ = 511.4$

Example 557

5



A.




10

A solution of 2-amino-5-cresol (5.0 g; 40 mmol), KOH (3.2 g; 57 mmol) was refluxed in EtOH (50 mL) and CS₂ (40 mL) for 8 h, after which the reaction mixture was concentrated in vacuo. The residue was partitioned between aq 1M HCl (100 mL) and EtOAc (200 mL). The organic phase was washed with water (2 x 100 mL), dried (MgSO₄) and concentrated in vacuo to give Part A compound (4.0 g; 60%) as a white powder.

20


B.



25

A solution of Part A compound (3.2 g; 19 mmol) and POCl₃ (3.75 g; 19 mmol) in toluene (150 mL) was heated at reflux for 2 h. The reaction mixture was washed successively with water and aqueous NaHCO₃, then dried,

5 c.



20


CC1=C(C(=O)O)N(C1Cc2ccc(OCC3=C(C)N(C3=O)C4=CC=CC=C4)cc2)Cc5cc6ncc(C)c6cc5

30

5

COc1ccc(cc1)Oc2cc(oc2C)c3ccc(OC)cc3C(=O)NCC(=O)OCC(Br)C(=O)CC(=O)OC

B.




A mixture of Part A compound (1.5 g, 7.2 mmol) and 4-methoxybenzamide (1.0 g, 6.6 mmol) was heated at 100°C for 2.5 h. The reaction mixture was chromatographed

(SiO₂; 5% acetone/CH₂Cl₂) to yield Part B compound (0.57 g, 33%).

C.


5



To a solution of the ester (0.57 g, 2.3 mmol) in THF (10 mL) was added LiAlH₄ (2.5 mL of a 1 M solution in THF, 2.5 mmol) dropwise over 10 min and the reaction was stirred at RT for 0.5 h. The reaction was quenched by adding a few drops of water and then partitioned between EtOAc (50 mL) and brine (10 mL). The organic phase was dried (MgSO₄) and concentrated in vacuo to give Part C (0.52 g, >95%) as an oil which was used in the following reaction without further purification.

D.


20



A mixture of Part C compound (0.52 g, 2.3 mmol), CH₃SO₂Cl (0.25 ml, 3.3 mmol) and Et₃N (0.5 ml, 3.6 mmol) in CH₂Cl₂ (10 mL) was stirred at RT for 12 h. Volatiles were removed in vacuo, and the residue was chromatographed (SiO₂; 4% acetone/CH₂Cl₂) to provide Part D compound (0.61 g, 85% for 2 steps) as a colorless oil.

E.

30



10

COc1ccc(cc1)C2=NC(=C(C)CC2OCC3=CC=CC=C3OC(=O)N(CCC(=O)OC(=O)C)OC4=CC=CC=C4OC)C5=CC=C(C=C5)OC

15

25


COC1=CC=C(C=C1)OC(=O)NCCOC2=CC=C(C=C2)OCC3=C(C)OC4=CC=C(C=C4)C5=CC(OC)=CC=C53

10

15

COC(=O)OCC#CC1=C(C)OC(=N1)c2ccccc2


25



[described in the synthesis of Example 503 Part B compound (50 mg; 0.13 mmol)], K₂CO₃ (17 mg; 0.34 mmol) in CH₃CN (1 mL) were heated at 70°C for 24 h. Additional K₂CO₃ (30 mg) and CH₃CN (1 mL) were added and the mixture was heated at 75°C for another 48 h. The reaction was cooled to RT, EtOAc was added, and the mixture was washed with aq 1M NaOH and brine. The organic phase was dried (Na₂SO₄) and concentrated in vacuo to give the crude product. This was purified by preparative HPLC (YMC S5 ODS 50 x 75 mm column; continuous gradient from 70:30 B:A to 100% B, where A = 90:10:0.1 H₂O:MeOH:TFA and B = 90:10:0.1 MeOH:H₂O:TFA) to give Part A compound (13 mg; 35%) as a colorless oil.

15


B.




To a solution of Part A compound (12 mg; 0.021 mmol) in 2:1 THF:H₂O (1.5 mL) was added LiOH (8 mg; 0.19 mmol). The solution was stirred at RT for 24 h, then acidified with excess 1M HCl (aq). The solution was extracted with EtOAc (2 x 5 mL). The combined organic extracts were washed with brine, dried (Na₂SO₄), and concentrated in vacuo. The crude product was purified by

$$[M + H]^+ = 541.3$$


Example 561

CCOC(=O)CN(C)C[C@H](c1ccc(OCC2=C(C)N(C2=O)c3ccccc3)cc1)COCC4=CC=CC=C4N5C(=C(C)C(=O)N5C6=CC=CC=C6)

A mixture of the mesylate (18 mg; 0061 mmol)




the phenol (50 mg; 0.13 mmol)



K₂CO₃ (17 mg; 0.34 mmol) in CH₃CN (1 mL) were heated at 70°C for 24 h. Additional K₂CO₃ (30 mg) and CH₃CN (1 mL) were added and the mixture was heated at 75°C for another 48 h. The reaction was cooled to RT, EtOAc was added, and the mixture was washed with aq 1M NaOH and brine.

The organic phase was dried (Na_2SO_4) and concentrated in vacuo to give the crude product. This was purified by preparative HPLC (YMC 55 ODS 50 x 75 mm column; continuous gradient from 70:30 B:A to 100% B, where A = 90:10:0.1 H_2O :MeOH:TFA and B = 90:10:0.1 MeOH: H_2O :TFA) to give Part A compound (13 mg; 35%) as a colorless oil.


B.



10


To a solution of Part A compound (12 mg; 0.021 mmol) in 2:1 THF: H_2O (1.5 mL) was added LiOH (8 mg; 0.19 mmol). The solution was stirred at RT for 24 h, then acidified with excess 1M HCl (aq). The solution was extracted with EtOAc (2 x 5 mL). The combined organic extracts were washed with brine, dried (Na_2SO_4), and concentrated in vacuo. The crude product was purified by preparative HPLC using the same conditions as above to give title compound. $[\text{M} + \text{H}]^+ = 541.3$

20


Example 562

25


A.




15 B.



30 C.



15 D.

CCOC1=CC=C(OC(=O)CN(Cc2ccc(C#CCc3c(C)c4c(c3)nc(c4)C5=CC=CC=C5)cc2)C1=CC=C5

10

Cc1c(C#CCc2ccc(cc2)CNC(=O)Oc3ccc(OC)cc3)nc(C4=CC=CC=C4)o1


15

25

Cc1c(CCCc2ccc(cc2)CN(C)C(=O)Oc3ccc(OC)cc3)c(O)c1c4ccccc4C(=O)O


30

A.




5

A solution of Example 562 Part E compound



(38 mg; 0.072 mmol) in MeOH (5 mL) was stirred under an atmosphere of H₂ in the presence of 10% Pd/C catalyst (10 mg) at RT for 2 h. The catalyst was filtered off and the filtrate was concentrated in vacuo to give Part A compound (35 mg; 92%) as an oil.

B.



15

A solution of Part A compound (35 mg; 0.066 mmol) in aqueous LiOH (1 mL of a 1M solution) and THF (5 mL) was stirred at RT for 2h. The reaction was acidified to pH 3 with excess aqueous 1M HCl and extracted with EtOAc (2 x 5 mL). The combined organic extracts were dried (Na₂SO₄) and concentrated in vacuo. The residue was purified by preparative HPLC (YMC S5 ODS reverse phase column; 30 x 250 mm; flow rate = 25 mL/min; 30 min continuous gradient from 50:50 A:B to 100% B, where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B =


Example 564



•




20


Example 565

5

A.




A mixture of Example 562 Part E compound




- 10 (80 mg; 0.15 mmol), quinoline (2 μ L; 0.01 mmol) and Lindlar's catalyst (7 mg; 5% Pd/CaCO₃) in toluene (2 mL) was stirred under an atmosphere of H₂ for 2 h. More Lindlar's catalyst (20 mg) was then added and stirring was continued under H₂ for another 2 h, after which the
- 15 reaction was complete by analytical HPLC. The reaction mixture was filtered (Celite®) and the filtrate was concentrated in vacuo. The residue was chromatographed (SiO₂; stepwise gradient from 3:1 to 2:1 hexane:EtOAc) to give Part A compound.

B.




- A solution of Part A compound and aqueous LiOH (1 mL
5 of a 1 M solution; 1 mmol) in THF was stirred at RT
overnight. The reaction mixture was acidified with
excess aqueous 1 M HCl and extracted with EtOAc (2x).
The combined organic extracts were concentrated in vacuo.
The residue was purified by preparative HPLC (as for
10 Example 495) to give the title compound (14 mg; 18%) as a
white solid. $[M + H]^+ = 513.3$


Example 566 (racemic)

15

A.




To a 0°C solution of Example 565 Part A compound



(60 mg; 0.11 mmol) in DCE (3 mL) was added dropwise
diethylzinc (43 μ L; 0.29 mmol). The solution was stirred
5 at 0°C for 10 min and iodochloromethane (244 μ L; 0.57
mmol) was then added. The reaction was allowed to warm
to RT and stirred at RT for 3 h, then was cautiously
quenched by addition of aqueous HCl (1 mL of a 1 M
solution). The aqueous layer was extracted with EtOAc
10 (2x); the combined organic extracts were dried (Na_2SO_4)
and concentrated in vacuo. The residue was
chromatographed (SiO_2 ; stepwise gradient from 3:1 to 2:1
hexane:EtOAc) to give crude Part A compound, which was
used in the next step without further purification.


15

B.




A solution of crude Part A compound and aqueous
20 LiOH (1 mL of a 1 M solution; 1 mmol) in THF was stirred
at RT overnight. The reaction mixture was acidified with
excess aqueous 1 M HCl and extracted with EtOAc. The
combined organic extracts were concentrated in vacuo.
The residue was purified by preparative HPLC (conditions)
25 to give the title compound (7 mg; 12% over 2 steps) as a
white solid. $[\text{M} + \text{H}]^+ = 527.2$.

Example 567




5

A.



A mixture of Example 562 Part D compound


10



15

(300 mg; 1.52 mmol), N-bromo-succinimide (297 mg; 1.67 mmol) and AgNO₃ (28 mg; 0.19 mmol) in acetone (2 mL) was stirred at RT for 30 min. The mixture was filtered and the filtrate was concentrated in vacuo. The residue was chromatographed (SiO₂; hexane:EtOAc 5:1) to give Part A compound (320 mg; 76%) as yellow crystals.


B.



20

To a solution of Part A compound (320 mg; 1.2 mmol), Ph_3P (13 mg; 0.05 mmol) and $\text{Tris}(\text{dibenzylidene-acetone})\text{dipalladium}(0)$ (5 mg; 0.006 mmol) in THF (1 mL) was added Bu_3SnH (700 μL ; 2.5 mmol) dropwise under an atmosphere of N_2 . The mixture was stirred at RT for 2 h, then was quenched by addition of aqueous KF (7 mL of a 1 M solution). The mixture was stirred vigorously overnight, then extracted with EtOAc (2x). The combined organic extracts were washed with H_2O , dried (Na_2SO_4) and


concentrated in vacuo. The residual oil was chromatographed (SiO_2 ; hexane:EtOAc 3:1) to give Part B compound (200 mg; 35%) as an oil. In addition, the byproduct vinyl compound



5


(100 mg; 43%) was also obtained.

C.



10


A solution of Part B compound (100 mg; 0.020 mmol) and Example 562 Part B compound



(100 mg; 0.22 mmol) and $(\text{Ph}_3\text{P})_4\text{Pd}^0$ (3 mg; 0.002 mmol) in toluene was heated at 100°C overnight under an atmosphere of N_2 . Volatiles were removed in vacuo and the residue was chromatographed (SiO_2 ; stepwise gradient from 3:1 to 2:1 hexane:EtOAc) to give Part C compound.

20

D.



A solution of crude Part C compound (in aqueous LiOH (1 mL of a 1M solution) and THF (5 mL) was stirred

25


10

Examples 568 to 572

15


Example No.	Structure	[M+H] ⁺
568		511.2
569		515.9
570		511.2
571		513.2
572		513.3

Example 573




5

A.




10 To a mixture of the amino-ester (27 mg; 0.073 mmol)



5-methyl-2-phenyl-thiazol-4-yl-ethanol (25 mg; 0.11 mmol; Maybridge) resin-bound Ph_3P (27 mg; 0.081 mmol) in CH_2Cl_2 (0.5 mL) was added DEAD (20 μL ; 0.13 mmol). The reaction was stirred at RT for 6 h, then was filtered. The filtrate was concentrated in vacuo and the residue was purified by preparative HPLC (YMC S5 ODS 30 x 100 mm column; flow rate = 50 mL/min; continuous gradient from 30:70 B:A to 100% B, where solvent A = 90:10:0.1 H_2O :MeOH:TFA and solvent B = 90:10:0.1 MeOH: H_2O :TFA) to furnish Part A compound.

B.




25

5

CC1=C(C2=CC=CC=C2N=S1)CCOC3=CC=C(C=C3)CN(CCC4=CC=C(C=C4)OC5=CC=CC=C5)C(=O)O


A.

CCOC(=O)N(Cc1ccc(O)cc1)Cc2ccc(Oc3ccccc3)cc2

5-methyl-2-phenyl-thiazol-4-yl-ethanol (25 mg; 0.11 mmol; Maybridge) resin-bound Ph_3P (32 mg; 0.096 mmol) in CH_2Cl_2 (0.5 mL) was added DEAD (20 μL ; 0.13 mmol). The reaction was stirred at RT for 6 h, then was filtered. The filtrate was concentrated in vacuo to give crude Part A compound.

Cc1nc(CCCOc2ccc(cc2)CN(CCOC(=O)O)c3ccc(Oc4ccccc4)cc3)c(c1)c5ccccc5

Example 575


CC(Br)C(=O)C=C

- 307 -

5

CC(C)C(=O)CCOc1ccc(cc1)CN(C)C(=O)Oc2ccc(OC)cc2COC(=O)C

A mixture of Example 559 Part E compound (0.60 g, 1.7 mmol),



15

CC1=C(CSC1c2cccnc2)CCOC3=CC=C(C=C3)COC(=O)NCC(=O)OC4=CC=C(OC)C=C4

10






Cc1c(CCCOc2ccc(cc2)C(=O)NCC(=O)Oc3ccc(OC)cc3)nn(c1)c4cccnc4

20



Examples 576 to 580

Following the procedures set out hereinbefore and in the working Examples, the following compounds were prepared.

5

Example No.	Structure	[M+H] ⁺
576		551.2; 553.2
577		547.2
578		531.2
579		535.2; 537.2
580		551.2; 553.2



Examples 581 and 582 were synthesized according to the general procedures described for Examples 313 and 314.

Example	Structure	[M+H] ⁺
581		499.2
582		499.1


5

Examples 583 and 584 were synthesized according to the general methods described hereinbefore (e.g. for Example 139) using 4-methoxy thiophenol.

10

Example	Structure	[M+H] ⁺
583		533.3
584		533.3


Example 584




5 ¹H NMR (CDCl₃; 400 MHz): δ 2.42 (s, 3H), 3.04 (br s; 2H), 3.79 (s, 3H), 4.03 (br s, 2H), 4.25 (br s, 2H), 4.70 (br s, 2H), 6.8-7.0 (m, 5H), 7.15-7.30 (m, 1H), 7.35-7.50 (m, 5H), 7.95-8.05 (m, 2H); 8.95 (br s, 1H)

10

Example 585




A.




15

To a -78°C solution of methyltriphenylphosphonium bromide (4.2 g; 11.8 mmol) in THF (60 mL) was added dropwise n-butyllithium (4.7 mL of a 2.5 M solution in hexane; 11.8 mmol). The solution was allowed to warm to RT and stirred at RT for 45 min. To this mixture was added dropwise a solution of the aldehyde (3.0 g; 9.8 mmol)



5

10



15

20

25

30

CC1=C(C2=CC=CC=C2O1)CCOC3=CC=C(C=C3)C[C@H](C4=CC=CC=C4)C(=O)OCC5=CC=CC=C5

5

To a 0°C solution of Part B compound (1.10 g; 2.33 mmol) in CH₂Cl₂ (12 mL) were successively added methanesulfonyl chloride (220 µL; 2.80 mmol) and Et₃N (420 µL; 3.03 mmol) dropwise. The reaction was stirred at 0°C for 2 h, then was partitioned between CH₂Cl₂ and aqueous 1N HCl. The organic phase was washed with water and brine, dried (MgSO₄) and concentrated in vacuo to provide Part D compound (1.10 g; 86%) as a solid.


15

CC1=C(C2=CC=CC=C2N=C3C(=C1)OC(=C3)C4=CC=CC=C4)CCOC5=CC=C(C=C5)C[C@H](C6=CC=CC=C6)C(=O)OC7=CC=CC=C7

A mixture of Part D compound (1.10 g; 2.0 mmol) and
20 LiBr (260 mg; 3.0 mmol) in acetone (4 mL) was heated at
50°C for 14 h. The reaction was then cooled to RT and
concentrated in vacuo. The residue was chromatographed
(SiO₂; stepwise gradient from 9:1 to 4:1 hex:EtOAc) to
give Part E compound (481 mg; 45%) as an oil.


25

F.



To a -78°C slurry of CuCN (17 mg; 0.19 mmol) in freshly distilled anhydrous THF (0.54 mL) was added dropwise n -butyllithium (150 μL of a 2.5 M solution in hexanes). The mixture was allowed to warm slowly to 0°C to generate the higher-order cuprate reagent as a clear tan solution. The reaction was then cooled to -50°C and a solution of Part E compound (50 mg; 0.094 mmol) in THF (0.4 mL) was added dropwise. The reaction was stirred at -50°C for 1h and then allowed to warm slowly to 0°C over 2h. The mixture was then quenched at 0°C by addition of 9:1 saturated aqueous NH_4Cl :concentrated NH_4OH (2 mL), and then allowed to warm to RT with vigorous stirring until complete dissolution had occurred. The aqueous phase was extracted with EtOAc (2x), and the combined organic extracts were washed with saturated aqueous NH_4Cl and brine, dried (MgSO_4) and concentrated in vacuo. The residue was chromatographed (SiO_2 ; stepwise gradient from 9:1 to 2:1 hex:EtOAc) to provide Part F compound (26 mg; 54%) as a solid.

G.




25

A mixture of Part F compound (26 mg; 0.051 mmol) and 10% palladium on carbon (10 mg) in 2:1 MeOH:EtOAc (1.2 mL) was stirred under an atmosphere of H₂ (balloon) at RT for 2 h, at which point the reaction was complete by HPLC. The catalyst was filtered off through Celite®

and the filtrate was concentrated in vacuo to give Part G compound (18 mg; 93%) as an oil.

H.




5

A solution of Part G compound (18 mg; 0.048 mmol), methyl bromoacetate (9 μ L; 0.095 mmol) and Et₃N (15 μ L; 0.10 mmol) in THF (500 μ L) was stirred at RT for 15h.

10 The reaction mixture was partitioned between H₂O and EtOAc (60 mL) each. The organic phase was washed with brine, dried (MgSO₄) and concentrated in vacuo to furnish crude Part H compound, which was used in the next step without further purification.

15


I.



A solution of crude Part H compound, 4-methoxyphenyl chloroformate (21 μ L; 0.143 mmol) and 4-dimethyl-aminopyridine (4 mg; 0.033 mmol) in pyridine (10 mL) was heated at 70°C for 2 h. The reaction mixture was partitioned between EtOAc and 1M aqueous HCl. The organic phase was washed with brine, dried (MgSO₄), and concentrated in vacuo. The residue was purified by preparative HPLC (YMC reverse phase ODS 20 x 100 mm column; 10 min continuous gradient from 50%A:50%B to 100%B + 10 min hold-time at 100% B, where Solvent A = 90:10:0.1 H₂O:MeOH:TFA, and Solvent B = 90:10:0.1 MeOH:H₂O:TFA; flow rate = 20 mL/min; retention time = 14.6

30

J.




To a solution of Part I compound (9.0 mg; 0.015 mmol) in THF:H₂O (750 µL of a 2:1 solution) was added LiOH.H₂O (2.5 mg; 0.06 mmol). The reaction was stirred at RT for 15 h; then EtOAc (2 mL) was added and the solution acidified with 1 N HCl solution to pH ~ 2. The organic phase was washed with water and brine, dried (MgSO₄) and concentrated in vacuo. The residue was purified by preparative HPLC (YMC reverse phase ODS 20 x 100 mm column; flow rate = 20 mL/min; 10 min continuous gradient from 50:50 B:A to 100% B + 10 min hold-time at 100% B, where solvent A = 90:10:0.1 H₂O:MeOH:TFA and solvent B = 90:10:0.1 MeOH:H₂O:TFA; retention time = 13.2 min) to provide the title compound (6.0 mg; 68%) as a white solid.

$$[M + H]^+ = 587.3$$


Example 586

25



The synthesis of Example 586 was performed using the identical sequence as described for Example 585 except

that the catalyst used in the aminohydroxylation procedure (step 2) for the preparation of the key intermediate




5

was hydroquinine 1,4-phthlazinediyl diether [(DHQ)₂PHAL; Aldrich] instead of hydroquinidine 1,4-phthlazinediyl diether [(DHQD)₂PHAL; Aldrich].


10

Example 587




15

A.




To a 0°C solution of ethyl propionylacetate (10.0 g, 69.4 mmol) in CHCl₃ (60 mL) was added dropwise a solution of Br₂ (3.6 mL; 69.4 mmol) in CHCl₃ (20 mL) and the resulting mixture was stirred at 0°C for 0.5 h. The reaction was allowed to warm to RT and stirred at RT for 0.5 h. Air was then bubbled into the mixture for 1 h. Volatiles were then removed in vacuo to yield an oily residue to provide crude Part A compound (15.3 g, >95% yield) as an oil which was used in the next reaction without further purification.

B.



To a mixture of piperonylic acid (2.0 g; 12 mmol), HOBT.H₂O (2.44 g; 18.1 mmol) and NH₄Cl (1.28 g; 23.7 mmol) in DMF (48 mL) were successively added EDCI.HCl (3.45 g; 18.1 mmol) and iPr₂NEt (2.3 mL; 48 mmol). The reaction mixture was stirred at RT overnight until the starting acid had been completely consumed (by HPLC). The mixture was partitioned between H₂O (80 mL) and EtOAc (250 mL). The aqueous phase was extracted with EtOAc (250 mL). The combined organic extracts were washed with aqueous 1 N HCl (40 mL), dried (Na₂SO₄) and concentrated in vacuo. The crude product was chromatographed (SiO₂; stepwise gradient from hex:EtOAc 1:1 to 100% EtOAc) to give Part B compound (1.5 g; 76 %) as a white solid.

C.




20

A mixture of Part A compound (2.11 g, 9.5 mmol) and Part B compound (1.41 g, 8.54 mmol) was heated with a heat gun until the mixture became homogeneous, after which the solution was heated at 130°C in an oil bath for 5 h. The reaction mixture was chromatographed (SiO₂; continuous gradient from hex to 4:1 Hex:EtOAc over 20 min, then continuous gradient from 4:1 Hex:EtOAc to 100% EtOAc over 15 min) to yield Part C compound (0.95 g, 39%) as a yellow solid.


30

D.




10

15




20

F.




30



and K_2CO_3 (15 mg, 0.109 mmol) in acetonitrile (1 mL) was shaken and heated at 80°C for 22 h. The reaction was cooled to RT and filtered. The filtrate was concentrated in vacuo and the residue was purified by preparative HPLC (YMC reverse-phase ODS 20 x 100 mm column; continuous gradient over 10 min from 70:30 A:B to 100% B, where A = 90:10:0.1 H_2O :MeOH:TFA, and B = 90:10:0.1 MeOH: H_2O :TFA, with 7 min hold time at 100% B; flow rate = 20 mL/min) to provide Part F compound (21 mg; 51%) as a colorless oil.

G.




15

A solution of Part F compound (21 mg, 0.037 mmol) and $LiOH \cdot H_2O$ (4.0 mg; 0.095 mmol) in THF- H_2O (2.0 mL of a 1:1 mixture) was shaken at RT for 4 h. The reaction mixture was acidified to pH 5 with 1 M aqueous HCl, then was extracted with EtOAc (3 mL) by shaking for 10 min. The organic phase was washed with H_2O (2 mL) and concentrated in vacuo to provide Example 586 (16.3 mg, 75%) as a solid foam.

$$[M + H]^+ = 561.2$$


25

Examples 588 to 596 were prepared according to the scheme described above.

Examples 588 to 591

Example No.	Ar	[M+H] ⁺
588		551.1
589		547.2
590		585.3
591		585.2

5

Examples 592 to 596

10

